



US008415512B2

(12) **United States Patent**
Sherman et al.

(10) **Patent No.:** **US 8,415,512 B2**
(45) **Date of Patent:** ***Apr. 9, 2013**

(54) **HYDROCARBON CONVERSION PROCESS IMPROVEMENTS**

(75) Inventors: **Jeffrey H. Sherman**, Vero Beach, FL (US); **Eric W. McFarland**, Santa Barbara, CA (US); **Michael J. Weiss**, Santa Barbara, CA (US); **Ivan Marc Lorkovic**, Santa Barbara, CA (US); **Leroy E. Laverman**, Santa Barbara, CA (US); **Shouli Sun**, Santa Barbara, CA (US); **Dieter J. Schaefer**, Goleta, CA (US); **Galen D. Stucky**, Santa Barbara, CA (US); **Peter C. Ford**, Santa Barbara, CA (US); **Philip Grosso**, Auburn, CA (US); **Ashley W. Reed**, Goleta, CA (US); **Michael F. Doherty**, Santa Barbara, CA (US)

(73) Assignee: **GRT, Inc.**, Santa Barbara, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **12/904,030**

(22) Filed: **Oct. 13, 2010**

(65) **Prior Publication Data**

US 2011/0034741 A1 Feb. 10, 2011

Related U.S. Application Data

(63) Continuation of application No. 12/692,831, filed on Jan. 25, 2010, now Pat. No. 7,838,708, which is a continuation of application No. 11/098,997, filed on Apr. 5, 2005, now abandoned, which is a continuation-in-part of application No. 10/430,240, filed on Aug. 19, 2003, now Pat. No. 7,161,050, which is a continuation-in-part of application No. 10/365,346, filed on Feb. 12, 2003, now abandoned, which is a continuation of application No. 10/298,440, filed on Nov. 19, 2002, now abandoned, which is a continuation-in-part of application No. 10/208,068, filed on Jul. 29, 2002, now abandoned, which is a continuation-in-part of application No. 10/054,004, filed on Jan. 24, 2002, now Pat. No. 6,486,368, which is a continuation-in-part of application No. 09/951,739, filed on Sep. 11, 2001, now Pat. No. 6,465,696, which is a continuation-in-part of application No. 09/886,078, filed on Jun. 20, 2001, now Pat. No. 6,472,572.

(51) **Int. Cl.**
C07C 27/00 (2006.01)
C07C 27/10 (2006.01)

(52) **U.S. Cl.**
USPC **568/893**; 568/890; 568/891; 570/223;
570/243; 570/245; 570/254

(58) **Field of Classification Search** 570/223,
570/243, 245, 254; 568/893, 891, 890

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,168,260 A 8/1939 Heisel et al.
2,246,082 A 6/1941 Vaughan et al.
2,488,083 A 11/1949 Gorin et al.
2,677,598 A 5/1954 Crummett et al.
2,941,014 A 6/1960 Rothweiler et al.
3,076,784 A 2/1963 Huermann et al.
3,172,915 A 3/1965 Borkowski et al.
3,246,043 A 4/1966 Rosset et al.
3,273,964 A 9/1966 Rosset
3,294,846 A 12/1966 Livak et al.
3,310,380 A 3/1967 Lester
3,346,340 A 10/1967 Louvar et al.
3,353,916 A 11/1967 Lester
3,353,919 A 11/1967 Stockman
3,496,242 A 2/1970 Berkowitz et al.
3,562,321 A 2/1971 Borkowski et al.
3,598,876 A 8/1971 Bloch
3,657,367 A 4/1972 Blake et al.
3,670,037 A 6/1972 Dugan
3,673,264 A 6/1972 Kuhn
3,679,758 A 7/1972 Schneider
3,702,886 A 11/1972 Argauer et al.
3,705,196 A 12/1972 Turner
3,799,997 A 3/1974 Schmerling
3,865,886 A 2/1975 Schindler et al.
3,876,715 A 4/1975 McNulty et al.
3,879,473 A 4/1975 Stapp

(Continued)

FOREIGN PATENT DOCUMENTS

BR 0210054 8/2004
CA 1099656 4/1981

(Continued)

OTHER PUBLICATIONS

U.S. Appl. No. 60/487,364, filed Jul. 15, 2003, Lorkovic et al.
U.S. Appl. No. 60/559,844, filed Apr. 6, 2004, Sherman et al.
U.S. Appl. No. 60/765,115, filed Feb. 3, 2006, Gadewar et al.
Abstract of JP2007045756, Hydrogenation method using diaphragm type hydrogenation catalyst, hydrogenation reaction apparatus and diaphragm type hydrogenation catalyst, Publication date: Feb. 22, 2007, Inventor: Shuji et al., esp@cenet database—worldwide.
Abstract of JP2007061594, Method for decomposing organohalogen compound and mobile decomposition system, Publication date: Mar. 15, 2007, Inventor: Koichi et al., esp@cenet database—worldwide.

(Continued)

Primary Examiner — Elvis O Price

(74) *Attorney, Agent, or Firm* — Adolph Locklar

(57) **ABSTRACT**

Improvements in previously disclosed methods of and apparatuses for converting alkanes, alkenes, and aromatics to olefins, alcohols, ethers, and aldehydes includes: safety improvements, use of alternative feedstocks, process simplification, improvements to the halogenation step, improvements to the reproporationation step, improvements to the solid oxide reaction, improvements to solid oxide regeneration, improvements in separations, maintenance, start-up, shut-down, and materials of construction.

20 Claims, No Drawings

U.S. PATENT DOCUMENTS							
3,879,480	A	4/1975	Riegel et al.	4,433,189	A	2/1984	Young
3,883,651	A	5/1975	Woitun et al.	4,433,192	A	2/1984	Olah
3,886,287	A	5/1975	Kobayashi et al.	4,439,409	A	3/1984	Puppe et al.
3,894,103	A	7/1975	Chang et al.	4,440,871	A	4/1984	Lok et al.
3,894,104	A	7/1975	Chang et al.	4,443,620	A	4/1984	Gelbein et al.
3,894,105	A	7/1975	Chang et al.	4,462,814	A	7/1984	Holmes et al.
3,894,107	A	7/1975	Butter et al.	4,465,884	A	8/1984	Degnan et al.
3,907,917	A	9/1975	Forth	4,465,893	A	8/1984	Olah
3,919,336	A	11/1975	Kurtz	4,467,130	A	8/1984	Olah
3,920,764	A	11/1975	Riegel et al.	4,467,133	A	8/1984	Chang et al.
3,923,913	A	12/1975	Antonini et al.	4,489,210	A	12/1984	Judat et al.
3,928,483	A	12/1975	Chang et al.	4,489,211	A	12/1984	Ogura et al.
3,965,205	A	6/1976	Garwood et al.	4,492,657	A	1/1985	Heiss
3,974,062	A	8/1976	Owen et al.	4,496,752	A	1/1985	Gelbein et al.
3,987,119	A	10/1976	Kurtz et al.	4,497,967	A	2/1985	Wan
3,992,466	A	11/1976	Plank et al.	4,499,314	A	2/1985	Seddon et al.
4,006,169	A	2/1977	Anderson et al.	4,506,105	A	3/1985	Kaufhold
4,011,278	A	3/1977	Plank et al.	4,509,955	A	4/1985	Hayashi
4,025,571	A	5/1977	Lago	4,513,092	A	4/1985	Chu et al.
4,025,572	A	5/1977	Lago	4,513,164	A	4/1985	Olah
4,025,575	A	5/1977	Chang et al.	4,523,040	A	6/1985	Olah
4,025,576	A	5/1977	Chang et al.	4,524,227	A	6/1985	Fowles et al.
4,035,285	A	7/1977	Owen et al.	4,524,228	A	6/1985	Fowles et al.
4,035,430	A	7/1977	Dwyer et al.	4,524,231	A	6/1985	Fowles et al.
4,039,600	A	8/1977	Chang	4,538,014	A	8/1985	Miale et al.
4,044,061	A	8/1977	Chang et al.	4,538,015	A	8/1985	Miale et al.
4,046,825	A	9/1977	Owen et al.	4,540,826	A	9/1985	Banasiak et al.
4,049,734	A	9/1977	Garwood et al.	4,543,434	A	9/1985	Chang
4,052,471	A	10/1977	Pearsall	4,544,781	A	10/1985	Chao et al.
4,052,472	A	10/1977	Givens et al.	4,547,612	A	10/1985	Tabak
4,058,576	A	11/1977	Chang et al.	4,550,217	A	10/1985	Graziani et al.
4,060,568	A	11/1977	Rodewald	4,550,218	A	10/1985	Chu
4,071,753	A	1/1978	Fulenwider et al.	4,568,660	A	2/1986	Klosiewicz
4,072,733	A	2/1978	Hargis et al.	4,579,977	A	4/1986	Drake
4,087,475	A	5/1978	Jordan	4,579,992	A	4/1986	Kaufhold et al.
4,088,706	A	5/1978	Kaeding	4,579,996	A	4/1986	Font Freide et al.
4,092,368	A	5/1978	Smith	4,587,375	A	5/1986	Debras et al.
4,110,180	A	8/1978	Nidola et al.	4,588,835	A	5/1986	Torii et al.
4,117,251	A	9/1978	Kaufhold et al.	4,590,310	A	5/1986	Townsend et al.
4,129,604	A	12/1978	Tsao	4,599,474	A	7/1986	Devries et al.
4,133,838	A	1/1979	Pearson	4,605,796	A	8/1986	Isogai et al.
4,133,966	A	1/1979	Pretzer et al.	4,605,803	A	8/1986	Chang et al.
4,138,440	A	2/1979	Chang et al.	4,621,161	A	11/1986	Shihabi
4,156,698	A	5/1979	Dwyer et al.	4,621,164	A	11/1986	Chang et al.
4,169,862	A	10/1979	Eden	4,633,027	A	12/1986	Owen et al.
4,172,099	A	10/1979	Severino	4,634,800	A	1/1987	Withers, Jr. et al.
4,187,255	A	2/1980	Dodd	4,642,403	A	2/1987	Hyde et al.
4,194,990	A	3/1980	Pieters et al.	4,642,404	A	2/1987	Shihabi
4,197,420	A	4/1980	Ferraris et al.	4,652,688	A	3/1987	Brophy et al.
4,219,680	A	8/1980	Konig et al.	4,654,449	A	3/1987	Chang et al.
4,249,031	A	2/1981	Drent et al.	4,655,893	A	4/1987	Beale
4,270,929	A	6/1981	Dang Vu et al.	4,658,073	A	4/1987	Tabak
4,272,338	A	6/1981	Lynch et al.	4,658,077	A	4/1987	Kolts et al.
4,282,159	A	8/1981	Davidson et al.	4,665,259	A	5/1987	Brazdil et al.
4,300,005	A	11/1981	Li	4,665,267	A	5/1987	Barri
4,300,009	A	11/1981	Haag et al.	4,665,270	A	5/1987	Brophy et al.
4,301,253	A	11/1981	Warren	4,675,410	A	6/1987	Feitler et al.
4,302,619	A	11/1981	Gross et al.	4,690,903	A	9/1987	Chen et al.
4,307,261	A	12/1981	Beard, Jr. et al.	4,695,663	A	9/1987	Hall et al.
4,308,403	A	12/1981	Knifton	4,696,985	A	9/1987	Martin
4,311,865	A	1/1982	Chen et al.	4,704,488	A	11/1987	Devries et al.
4,317,800	A	3/1982	Sloterdijk et al.	4,704,493	A	11/1987	Devries et al.
4,317,934	A	3/1982	Seemuth	4,709,108	A	11/1987	Devries et al.
4,317,943	A	3/1982	Knifton	4,720,600	A	1/1988	Beech, Jr. et al.
4,320,241	A	3/1982	Frankiewicz	4,720,602	A	1/1988	Chu
4,333,852	A	6/1982	Warren	4,724,275	A	2/1988	Hinnenkamp et al.
4,347,391	A	8/1982	Campbell	4,735,747	A	4/1988	Ollivier et al.
4,350,511	A	9/1982	Holmes et al.	4,737,594	A	4/1988	Olah
4,371,716	A	2/1983	Paxson et al.	4,748,013	A	5/1988	Saito et al.
4,373,109	A	2/1983	Olah	4,769,504	A	9/1988	Noceti et al.
4,376,019	A	3/1983	Gamlen et al.	4,774,216	A	9/1988	Kolts et al.
4,380,682	A	4/1983	Leitert et al.	4,775,462	A	10/1988	Imai et al.
4,384,159	A	5/1983	Diesen	4,777,321	A	10/1988	Harandi et al.
4,389,391	A	6/1983	Dunn	4,781,733	A	11/1988	Babcock et al.
4,410,714	A	10/1983	Apanel	4,783,566	A	11/1988	Kocal et al.
4,412,086	A	10/1983	Beard, Jr. et al.	4,788,369	A	11/1988	Marsh et al.
4,418,236	A	11/1983	Cornelius et al.	4,788,377	A	11/1988	Chang et al.
4,431,856	A	2/1984	Daviduk et al.	4,792,642	A	12/1988	Rule et al.
				4,795,732	A	1/1989	Bam

US 8,415,512 B2

4,795,737 A	1/1989	Rule et al.	5,157,189 A	10/1992	Karra
4,795,843 A	1/1989	Imai et al.	5,160,502 A	11/1992	Kimble et al.
4,795,848 A	1/1989	Teller et al.	5,166,452 A	11/1992	Gradl et al.
4,804,797 A	2/1989	Minet et al.	5,175,382 A	12/1992	Hebgen et al.
4,804,800 A	2/1989	Bortinger et al.	5,178,748 A	1/1993	Casci et al.
4,808,763 A	2/1989	Shum	5,185,479 A	2/1993	Stauffer
4,814,527 A	3/1989	Diesen	5,188,725 A	2/1993	Harandi
4,814,532 A	3/1989	Yoshida et al.	5,191,142 A	3/1993	Marshall et al.
4,814,535 A	3/1989	Yurchak	5,194,244 A	3/1993	Brownscombe et al.
4,814,536 A	3/1989	Yurchak	5,202,506 A	4/1993	Kirchner et al.
4,849,562 A	7/1989	Buhs et al.	5,202,511 A	4/1993	Salinas, III et al.
4,849,573 A	7/1989	Kaefing	5,210,357 A	5/1993	Kolts et al.
4,851,602 A	7/1989	Harandi et al.	5,215,648 A	6/1993	Zones et al.
4,851,606 A	7/1989	Ragonese et al.	5,223,471 A	6/1993	Washecheck
4,886,925 A	12/1989	Harandi	5,228,888 A	7/1993	Gmelin et al.
4,886,932 A	12/1989	Leyshon	5,233,113 A	8/1993	Periana et al.
4,891,463 A	1/1990	Chu	5,237,115 A	8/1993	Makovec et al.
4,895,995 A	1/1990	James, Jr. et al.	5,243,098 A	9/1993	Miller et al.
4,899,000 A	2/1990	Stauffer	5,243,114 A	9/1993	Johnson et al.
4,899,001 A	2/1990	Kalnes et al.	5,245,109 A	9/1993	Kaminsky et al.
4,899,002 A	2/1990	Harandi et al.	5,254,772 A	10/1993	Dukat et al.
4,902,842 A	2/1990	Kalnes et al.	5,254,790 A	10/1993	Thomas et al.
4,925,995 A	5/1990	Robschlager	5,264,635 A	11/1993	Le et al.
4,929,781 A	5/1990	James, Jr. et al.	5,268,518 A	12/1993	West et al.
4,939,310 A	7/1990	Wade	5,276,226 A	1/1994	Horvath et al.
4,939,311 A	7/1990	Washecheck et al.	5,276,240 A	1/1994	Timmons et al.
4,945,175 A	7/1990	Hobbs et al.	5,276,242 A	1/1994	Wu
4,950,811 A	8/1990	Doussain et al.	5,284,990 A	2/1994	Peterson et al.
4,950,822 A	8/1990	Dileo et al.	5,300,126 A	4/1994	Brown et al.
4,956,521 A	9/1990	Volles	5,306,855 A	4/1994	Periana et al.
4,962,252 A	10/1990	Wade	5,316,995 A	5/1994	Kaminsky et al.
4,973,776 A	11/1990	Allenger et al.	5,319,132 A	6/1994	Ozawa et al.
4,973,786 A	11/1990	Karra	5,334,777 A	8/1994	Miller et al.
4,982,024 A	1/1991	Lin et al.	5,345,021 A	9/1994	Casci et al.
4,982,041 A	1/1991	Campbell	5,354,916 A	10/1994	Horvath et al.
4,988,660 A	1/1991	Campbell	5,354,931 A	10/1994	Jan et al.
4,990,696 A	2/1991	Stauffer	5,366,949 A	11/1994	Schubert
4,990,711 A	2/1991	Chen et al.	5,371,313 A	12/1994	Ostrowicki
5,001,293 A	3/1991	Nubel et al.	5,382,704 A	1/1995	Krespan et al.
5,004,847 A	4/1991	Beaver et al.	5,382,743 A	1/1995	Beech, Jr. et al.
5,013,424 A	5/1991	James, Jr. et al.	5,382,744 A	1/1995	Abbott et al.
5,013,793 A	5/1991	Wang et al.	5,385,718 A	1/1995	Casci et al.
5,019,652 A	5/1991	Taylor et al.	5,395,981 A	3/1995	Marker
5,026,934 A	6/1991	Bains et al.	5,399,258 A	3/1995	Fletcher et al.
5,026,937 A	6/1991	Bricker	5,401,890 A	3/1995	Parks
5,026,944 A	6/1991	Allenger et al.	5,401,894 A	3/1995	Brasier et al.
5,034,566 A	7/1991	Ishino et al.	5,406,017 A	4/1995	Withers, Jr.
5,043,502 A	8/1991	Martindale et al.	5,414,173 A	5/1995	Garces et al.
5,055,235 A	10/1991	Brackenridge et al.	5,430,210 A	7/1995	Grasselli et al.
5,055,633 A	10/1991	Volles	5,430,214 A	7/1995	Smith et al.
5,055,634 A	10/1991	Volles	5,430,219 A	7/1995	Sanfilippo et al.
5,059,744 A	10/1991	Harandi et al.	5,436,378 A	7/1995	Masini et al.
5,068,478 A	11/1991	Miller et al.	5,444,168 A	8/1995	Brown
5,071,449 A	12/1991	Sircar	5,446,234 A	8/1995	Casci et al.
5,071,815 A	12/1991	Wallace	5,453,557 A	9/1995	Harley et al.
5,073,656 A	12/1991	Chafin et al.	5,456,822 A	10/1995	Marcilly et al.
5,073,657 A	12/1991	Warren	5,457,255 A	10/1995	Kumata et al.
5,082,473 A	1/1992	Keefer	5,464,799 A	11/1995	Casci et al.
5,082,816 A	1/1992	Teller et al.	5,465,699 A	11/1995	Voigt
5,085,674 A	2/1992	Leavitt	5,470,377 A	11/1995	Whitlock
5,087,779 A	2/1992	Nubel et al.	5,480,629 A	1/1996	Thompson et al.
5,087,786 A	2/1992	Nubel et al.	5,486,627 A	1/1996	Quarderer et al.
5,087,787 A	2/1992	Kimble et al.	5,489,719 A	2/1996	Le et al.
5,093,542 A	3/1992	Gaffney	5,489,727 A	2/1996	Randolph et al.
5,096,469 A	3/1992	Keefer	5,500,297 A	3/1996	Thompson et al.
5,097,083 A	3/1992	Stauffer	5,510,525 A	4/1996	Sen et al.
5,099,084 A	3/1992	Stauffer	5,523,503 A	6/1996	Funk et al.
5,105,045 A	4/1992	Kimble et al.	5,525,230 A	6/1996	Wrigley et al.
5,105,046 A	4/1992	Washecheck	5,538,540 A	7/1996	Whitlock
5,107,032 A	4/1992	Erb et al.	5,563,313 A	10/1996	Chung et al.
5,107,051 A	4/1992	Pannell	5,565,092 A	10/1996	Pannell et al.
5,107,061 A	4/1992	Ou et al.	5,565,616 A	10/1996	Li et al.
5,108,579 A	4/1992	Casci	5,571,762 A	11/1996	Clerici et al.
5,118,899 A	6/1992	Kimble et al.	5,571,885 A	11/1996	Chung et al.
5,120,332 A	6/1992	Wells	5,599,381 A	2/1997	Whitlock
5,132,343 A	7/1992	Zwecker et al.	5,600,043 A	2/1997	Johnston et al.
5,138,112 A	8/1992	Gosling et al.	5,600,045 A	2/1997	Van Der Aalst et al.
5,139,991 A	8/1992	Taylor et al.	5,609,654 A	3/1997	Le et al.
5,146,027 A	9/1992	Gaffney	5,633,419 A	5/1997	Spencer et al.

US 8,415,512 B2

5,639,930 A	6/1997	Penick	6,320,085 B1	11/2001	Arvai et al.
5,653,956 A	8/1997	Zones	6,337,063 B1	1/2002	Rouleau et al.
5,656,149 A	8/1997	Zones et al.	6,342,200 B1	1/2002	Rouleau et al.
5,661,097 A	8/1997	Spencer et al.	6,368,490 B1	4/2002	Gestermann
5,663,465 A	9/1997	Clegg et al.	6,369,283 B1	4/2002	Guram et al.
5,663,474 A	9/1997	Pham et al.	6,372,949 B1	4/2002	Brown et al.
5,675,046 A	10/1997	Ohno et al.	6,376,731 B1	4/2002	Evans et al.
5,675,052 A	10/1997	Menon et al.	6,380,328 B1	4/2002	McConville et al.
5,679,134 A	10/1997	Brugerolle et al.	6,380,423 B2	4/2002	Banning et al.
5,679,879 A	10/1997	Mercier et al.	6,380,444 B1	4/2002	Bjerrum et al.
5,684,213 A	11/1997	Nemphos et al.	6,395,945 B1	5/2002	Randolph
5,693,191 A	12/1997	Pividal et al.	6,403,840 B1	6/2002	Zhou et al.
5,695,890 A	12/1997	Thompson et al.	6,406,523 B1	6/2002	Connor et al.
5,698,747 A	12/1997	Godwin et al.	6,423,211 B1	7/2002	Randolph et al.
5,705,712 A	1/1998	Frey et al.	6,426,441 B1	7/2002	Randolph et al.
5,705,728 A	1/1998	Viswanathan et al.	6,426,442 B1	7/2002	Ichikawa et al.
5,705,729 A	1/1998	Huang	6,452,058 B1 *	9/2002	Schweizer et al. 570/223
5,708,246 A	1/1998	Camaioni et al.	6,455,650 B1	9/2002	Lipian et al.
5,720,858 A	2/1998	Noceti et al.	6,462,243 B1	10/2002	Zhou et al.
5,728,897 A	3/1998	Buysch et al.	6,465,699 B1	10/2002	Grosso
5,728,905 A	3/1998	Clegg et al.	6,465,969 B1	10/2002	Zhou et al.
5,734,073 A	3/1998	Chambers et al.	6,472,345 B2	10/2002	Hintermann et al.
5,741,949 A	4/1998	Mack	6,472,572 B1	10/2002	Zhou et al.
5,744,669 A	4/1998	Kalnes et al.	6,475,463 B1	11/2002	Elomari et al.
5,750,801 A	5/1998	Buysch et al.	6,475,464 B1	11/2002	Rouleau et al.
5,770,175 A	6/1998	Zones	6,479,705 B2	11/2002	Murata et al.
5,776,871 A	7/1998	Cothran et al.	6,482,997 B2	11/2002	Petit-Clair et al.
5,780,703 A	7/1998	Chang et al.	6,486,368 B1	11/2002	Zhou et al.
5,798,314 A	8/1998	Spencer et al.	6,495,484 B1	12/2002	Holtcamp
5,814,715 A	9/1998	Chen et al.	6,509,485 B2	1/2003	Mul et al.
5,817,904 A	10/1998	Vic et al.	6,511,526 B2	1/2003	Jagger et al.
5,821,394 A	10/1998	Schoebrechts et al.	6,514,319 B2	2/2003	Keefer et al.
5,847,224 A	12/1998	Koga et al.	6,518,474 B1	2/2003	Sanderson et al.
5,849,978 A	12/1998	Benazzi et al.	6,518,476 B1	2/2003	Culp et al.
5,866,735 A	2/1999	Cheung et al.	6,525,228 B2	2/2003	Chauvin et al.
5,895,831 A	4/1999	Brasier et al.	6,525,230 B2	2/2003	Grosso
5,898,086 A	4/1999	Harris	6,528,693 B1	3/2003	Gandy et al.
5,905,169 A	5/1999	Jacobson	6,538,162 B2	3/2003	Chang et al.
5,906,892 A	5/1999	Thompson et al.	6,540,905 B1	4/2003	Elomari
5,908,963 A	6/1999	Voss et al.	6,545,191 B1	4/2003	Stauffer
5,952,538 A	9/1999	Vaughn et al.	6,547,958 B1	4/2003	Elomari
5,959,170 A	9/1999	Withers	6,548,040 B1	4/2003	Rouleau et al.
5,968,236 A	10/1999	Bassine	6,552,241 B1	4/2003	Randolph et al.
5,969,195 A	10/1999	Stabel et al.	6,566,572 B2	5/2003	Okamoto et al.
5,977,402 A	11/1999	Sekiguchi et al.	6,572,829 B2	6/2003	Linkous et al.
5,983,476 A	11/1999	Eshelman et al.	6,585,953 B2	7/2003	Roberts et al.
5,986,158 A	11/1999	Van Broekhoven et al.	6,616,830 B2	9/2003	Elomari
5,994,604 A	11/1999	Reagen et al.	6,620,757 B2	9/2003	McConville et al.
5,998,679 A	12/1999	Miller	6,632,971 B2	10/2003	Brown et al.
5,998,686 A	12/1999	Clem et al.	6,635,793 B2	10/2003	Mul et al.
6,002,059 A	12/1999	Hellring et al.	6,641,644 B2	11/2003	Jagger et al.
6,015,867 A	1/2000	Fushimi et al.	6,646,102 B2	11/2003	Boriack et al.
6,018,088 A	1/2000	Olah	6,669,846 B2	12/2003	Perriello
6,022,929 A	2/2000	Chen et al.	6,672,572 B2	1/2004	Werlen
6,034,288 A	3/2000	Scott et al.	6,679,986 B1	1/2004	Da Silva et al.
6,056,804 A	5/2000	Keefer et al.	6,680,415 B1	1/2004	Gulotty, Jr. et al.
6,068,679 A	5/2000	Zheng	6,692,626 B2	2/2004	Keefer et al.
6,072,091 A	6/2000	Cosyns et al.	6,692,723 B2	2/2004	Rouleau et al.
6,087,294 A	7/2000	Klabunde et al.	6,710,213 B2	3/2004	Aoki et al.
6,090,312 A	7/2000	Ziaka et al.	6,713,087 B2	3/2004	Tracy et al.
6,096,932 A	8/2000	Subramanian	6,713,655 B2	3/2004	Yilmaz et al.
6,096,933 A	8/2000	Cheung et al.	RE38,493 E	4/2004	Keefer et al.
6,103,215 A	8/2000	Zones et al.	6,723,808 B2	4/2004	Holtcamp
6,107,561 A	8/2000	Thompson	6,727,400 B2	4/2004	Messier et al.
6,117,371 A	9/2000	Mack	6,740,146 B2	5/2004	Simonds
6,124,514 A	9/2000	Emmrich et al.	6,753,390 B2	6/2004	Ehrman et al.
6,127,588 A	10/2000	Kimble et al.	6,765,120 B2	7/2004	Weber et al.
6,130,260 A	10/2000	Hall et al.	6,797,845 B1	9/2004	Hickman et al.
6,143,939 A	11/2000	Farcasiu et al.	6,797,851 B2	9/2004	Martens et al.
6,169,218 B1	1/2001	Hearn et al.	6,821,924 B2	11/2004	Gulotty, Jr. et al.
6,180,841 B1	1/2001	Fatutto et al.	6,822,123 B2	11/2004	Stauffer
6,187,871 B1	2/2001	Thompson et al.	6,822,125 B2	11/2004	Lee et al.
6,187,983 B1	2/2001	Sun	6,825,307 B2	11/2004	Goodall
6,203,712 B1	3/2001	Bronner et al.	6,825,383 B1	11/2004	Dewkar et al.
6,207,864 B1	3/2001	Henningsen et al.	6,831,032 B2	12/2004	Spaether
6,225,517 B1	5/2001	Nascimento et al.	6,838,576 B1	1/2005	Wicki et al.
6,248,218 B1	6/2001	Linkous et al.	6,841,063 B2	1/2005	Elomari
6,265,505 B1	7/2001	McConville et al.	6,852,896 B2	2/2005	Stauffer
6,281,405 B1	8/2001	Davis et al.	6,866,950 B2	3/2005	Connor et al.

6,869,903 B2	3/2005	Matsunaga	7,348,295 B2	3/2008	Zones et al.
6,875,339 B2	4/2005	Rangarajan et al.	7,348,464 B2	3/2008	Waycuilis
6,878,853 B2	4/2005	Tanaka et al.	7,357,904 B2	4/2008	Zones et al.
6,888,013 B2	5/2005	Paparatto et al.	7,361,794 B2	4/2008	Grosso
6,900,363 B2	5/2005	Harth et al.	7,390,395 B2	6/2008	Elomari
6,902,602 B2	6/2005	Keefe et al.	2002/0102672 A1	8/2002	Mizrahi
6,903,171 B2	6/2005	Rhodes et al.	2002/0198416 A1	12/2002	Zhou et al.
6,909,024 B1	6/2005	Jones et al.	2003/0004380 A1	1/2003	Grumann
6,921,597 B2	7/2005	Keefe et al.	2003/0065239 A1	4/2003	Zhu
6,933,417 B1	8/2005	Henley et al.	2003/0069452 A1	4/2003	Sherman et al.
6,946,566 B2	9/2005	Yaegashi et al.	2003/0078456 A1	4/2003	Yilmaz et al.
6,953,868 B2	10/2005	Boaen et al.	2003/0120121 A1	6/2003	Sherman et al.
6,953,873 B2	10/2005	Cortright et al.	2003/0125589 A1	7/2003	Grosso
6,956,140 B2	10/2005	Ehrenfeld	2003/0166973 A1	9/2003	Zhou et al.
6,958,306 B2	10/2005	Holtcamp	2004/0006246 A1	1/2004	Sherman et al.
6,984,763 B2	1/2006	Schweizer et al.	2004/0062705 A1	4/2004	Leduc
7,001,872 B2	2/2006	Pyecroft et al.	2004/0152929 A1	8/2004	Clarke
7,002,050 B2	2/2006	Santiago Fernandez et al.	2004/0158107 A1	8/2004	Aoki
7,011,811 B2	3/2006	Elomari	2004/0158108 A1	8/2004	Snoble
7,019,182 B2	3/2006	Grosso	2004/0187684 A1	9/2004	Elomari
7,026,145 B2	4/2006	Mizrahi et al.	2005/0047927 A1	3/2005	Lee et al.
7,026,519 B2	4/2006	Santiago Fernandez et al.	2005/0148805 A1	7/2005	Jones
7,037,358 B2	5/2006	Babicki et al.	2005/0171393 A1	8/2005	Lorkovic
7,045,670 B2	5/2006	Johnson et al.	2005/0192468 A1	9/2005	Sherman et al.
7,049,388 B2	5/2006	Boriack et al.	2005/0215837 A1	9/2005	Hoffpauir
7,053,252 B2	5/2006	Boussand et al.	2005/0234276 A1	10/2005	Waycuilis
7,057,081 B2	6/2006	Allison et al.	2005/0245772 A1	11/2005	Fong
7,060,865 B2	6/2006	Ding et al.	2005/0245777 A1	11/2005	Fong
7,064,238 B2	6/2006	Waycuilis	2005/0267224 A1	12/2005	Herling
7,064,240 B2	6/2006	Ohno et al.	2006/0025617 A1	2/2006	Begley
7,067,448 B1	6/2006	Weitkamp et al.	2006/0100469 A1	5/2006	Waycuilis
7,083,714 B2	8/2006	Elomari	2006/0135823 A1	6/2006	Jun
7,084,308 B1	8/2006	Stauffer	2006/0138025 A1	6/2006	Zones
7,091,270 B2	8/2006	Zilberman et al.	2006/0138026 A1	6/2006	Chen
7,091,387 B2	8/2006	Fong et al.	2006/0149116 A1	7/2006	Slaugh
7,091,391 B2	8/2006	Stauffer	2006/0229228 A1	10/2006	Komon et al.
7,094,936 B1	8/2006	Owens et al.	2006/0229475 A1	10/2006	Weiss et al.
7,098,371 B2	8/2006	Mack et al.	2006/0270863 A1	11/2006	Reiling
7,105,710 B2	9/2006	Boons et al.	2006/0288690 A1	12/2006	Elomari
7,138,534 B2	11/2006	Forlin et al.	2007/0004955 A1	1/2007	Kay
7,141,708 B2	11/2006	Marsella et al.	2007/0078285 A1	4/2007	Dagle
7,145,045 B2	12/2006	Harmsen et al.	2007/0100189 A1	5/2007	Stauffer
7,148,356 B2	12/2006	Smith, III et al.	2007/0129584 A1	6/2007	Basset
7,148,390 B2	12/2006	Zhou et al.	2007/0142680 A1	6/2007	Ayoub
7,151,199 B2	12/2006	Martens et al.	2007/0148067 A1	6/2007	Zones
7,161,050 B2	1/2007	Sherman et al.	2007/0148086 A1	6/2007	Zones
7,169,730 B2	1/2007	Ma et al.	2007/0149778 A1	6/2007	Zones
7,176,340 B2	2/2007	Van Broekhoven et al.	2007/0149789 A1	6/2007	Zones
7,176,342 B2	2/2007	Bellussi et al.	2007/0149819 A1	6/2007	Zones
7,182,871 B2	2/2007	Perriello	2007/0149824 A1	6/2007	Zones
7,193,093 B2	3/2007	Murray et al.	2007/0149837 A1	6/2007	Zones
7,196,239 B2	3/2007	Van Egmond et al.	2007/0197801 A1	8/2007	Bolk
7,199,083 B2	4/2007	Zevallos	2007/0197847 A1	8/2007	Liu
7,199,255 B2	4/2007	Murray et al.	2007/0213545 A1	9/2007	Bolk
7,208,641 B2	4/2007	Nagasaki et al.	2007/0238905 A1	10/2007	Arredondo
7,214,750 B2	5/2007	McDonald et al.	2007/0238909 A1	10/2007	Gadewar et al.
7,220,391 B1	5/2007	Huang et al.	2007/0251382 A1	11/2007	Gadewar
7,226,569 B2	6/2007	Elomari	2007/0276168 A1	11/2007	Garel
7,226,576 B2	6/2007	Elomari	2007/0284284 A1	12/2007	Zones
7,230,150 B2	6/2007	Grosso et al.	2008/0171898 A1	7/2008	Waycuilis
7,230,151 B2	6/2007	Martens et al.	2008/0183022 A1	7/2008	Waycuilis
7,232,872 B2	6/2007	Shaffer et al.	2008/0188697 A1	8/2008	Lorkovic
7,238,846 B2	7/2007	Janssen et al.	2008/0269534 A1	10/2008	Lorkovic
7,244,795 B2	7/2007	Agapiou et al.	2008/0314758 A1	12/2008	Grosso
7,244,867 B2	7/2007	Waycuilis	2009/0069606 A1	3/2009	Komon
7,250,107 B2	7/2007	Benazzi et al.	2009/0127156 A1	5/2009	Tung
7,250,542 B2	7/2007	Smith, Jr. et al.	2010/0096588 A1	4/2010	Gadewar
7,252,920 B2	8/2007	Kurokawa et al.	2010/0099928 A1	4/2010	Gadewar
7,253,327 B2	8/2007	Janssens et al.	2010/0099929 A1	4/2010	Gadewar
7,253,328 B2	8/2007	Stauffer	2010/0099930 A1	4/2010	Stoimenov
7,265,193 B2	9/2007	Weng et al.	2010/0105972 A1	4/2010	Lorkovic
7,267,758 B2	9/2007	Benazzi et al.	2010/0121119 A1	5/2010	Sherman
7,268,263 B1	9/2007	Frey et al.			
7,271,303 B1	9/2007	Sechrist et al.			
7,273,957 B2	9/2007	Bakshi et al.			
7,282,603 B2	10/2007	Richards	CA	1101441	5/1981
7,285,698 B2	10/2007	Liu et al.	CA	1202610	4/1986
7,304,193 B1	12/2007	Frey et al.	CA	2447761 A1	11/2002
7,342,144 B2	3/2008	Kaizik et al.	CA	2471295 A1	7/2003
			CA	2542857	5/2005

FOREIGN PATENT DOCUMENTS

US 8,415,512 B2

CA	2236126	8/2006	WO	01/14300	A1	3/2001
CA	2203115	9/2006	WO	01/38275	A1	5/2001
CA	2510093	12/2006	WO	01/44149	A1	6/2001
EP	0021497	1/1981	WO	02/094749	A1	11/2002
EP	0164798	A1 12/1985	WO	02/094750	A1	11/2002
EP	0418971	A1 3/1991	WO	02/094751	A2	11/2002
EP	0418974	A1 3/1991	WO	02/094752	A1	11/2002
EP	0418975	A1 3/1991	WO	03/000635	A1	1/2003
EP	0510238	A1 10/1992	WO	03/002251	A2	1/2003
EP	0526908	A2 2/1993	WO	03/018524	A1	3/2003
EP	0346612	B1 8/1993	WO	03/020676	A1	3/2003
EP	0560546	A1 9/1993	WO	03/022827	A1	3/2003
EP	0976705	A1 7/1998	WO	03/043575	A2	5/2003
EP	1186591	A2 3/2002	WO	03/051813	A1	6/2003
EP	1253126	A1 10/2002	WO	03/062143	A1	7/2003
EP	1312411	A2 5/2003	WO	03/062172	A2	7/2003
EP	1395536	3/2004	WO	03/078366	A1	9/2003
EP	1404636	4/2004	WO	2004/018093	A2	3/2004
EP	1235769	B1 5/2004	WO	2004/067487	A2	8/2004
EP	1435349	A2 7/2004	WO	2005/014168	A1	2/2005
EP	1440939	A1 7/2004	WO	2005/019143	A1	3/2005
EP	1474371	11/2004	WO	2005/021468	A1	3/2005
EP	1235772	B1 1/2005	WO	2005/035121	A2	4/2005
EP	1661620	A1 5/2006	WO	2005/037758	A1	4/2005
EP	1760057	A1 3/2007	WO	2005/054120	A2	6/2005
EP	1689728	B1 4/2007	WO	2005/056525	A2	6/2005
EP	1808227	A1 7/2007	WO	2005/058782	A1	6/2005
EP	1837320	A1 9/2007	WO	2005/090272	A1	9/2005
GB	5125	0/1912	WO	2005/095310	A2	10/2005
GB	156122	3/1922	WO	2005/105709	A1	11/2005
GB	294100	6/1929	WO	2005/105715	A1	11/2005
GB	363009	12/1931	WO	2005/110953	A1	11/2005
GB	402928	12/1933	WO	2005/113437	A1	12/2005
GB	474922	A 11/1937	WO	2005/113440	A1	12/2005
GB	536491	5/1941	WO	2006/007093	A1	1/2006
GB	553950	6/1943	WO	2006/015824	A1	2/2006
GB	586483	3/1947	WO	2006/019399	A2	2/2006
GB	775590	5/1957	WO	2006/020234	A1	2/2006
GB	793214	4/1958	WO	2006/036293	A1	4/2006
GB	796048	6/1958	WO	2006/039213	A1	4/2006
GB	796085	6/1958	WO	2006/039354	A2	4/2006
GB	883256	11/1961	WO	2006/043075	A1	4/2006
GB	950975	3/1964	WO	2006/053345	A1	5/2006
GB	950976	3/1964	WO	2006-067155	A2	6/2006
GB	991303	5/1965	WO	2006/067188	A1	6/2006
GB	995960	6/1965	WO	2006/067190	A1	6/2006
GB	1015033	12/1965	WO	2006/067191	A1	6/2006
GB	1104294	2/1968	WO	2006/067192	A1	6/2006
GB	1133752	11/1968	WO	2006/067193	A1	6/2006
GB	1172002	11/1969	WO	2006/069107	A2	6/2006
GB	1212240	11/1970	WO	2006/071354	A1	7/2006
GB	1233299	5/1971	WO	2006/076942	A1	7/2006
GB	1253618	11/1971	WO	2006/083427	A1	8/2006
GB	1263806	A 2/1972	WO	2006-100312	A2	9/2006
GB	1446803	8/1976	WO	2006/104909	A2	10/2006
GB	1542112	3/1979	WO	2006/104914	A1	10/2006
GB	2095243	A 9/1982	WO	2006/111997	A1	10/2006
GB	2095245	A 9/1982	WO	2006/113205	A2	10/2006
GB	2095249	A 9/1982	WO	2006/118935	A2	11/2006
GB	2116546	A 9/1982	WO	2007/001934	A2	1/2007
GB	2120249	A 11/1983	WO	2007/017900	A2	2/2007
GB	2185754	A 7/1987	WO	2007/044139	A1	4/2007
GB	2191214	A 12/1987	WO	2007/046986	A2	4/2007
JP	2004-529189	9/2004	WO	2007/050745	A1	5/2007
WO	83/00859	3/1983	WO	2007/071046	A1	6/2007
WO	85/04863	11/1985	WO	2007/079038	A2	7/2007
WO	90/08120	7/1990	WO	2007/091009	A2	8/2007
WO	90/08752	8/1990	WO	2007/094995	A2	8/2007
WO	91/18856	12/1991	WO	2007/107031	A1	9/2007
WO	92/03401	3/1992	WO	2007/111997	A2	10/2007
WO	92/12946	8/1992	WO	2007/114479	A1	10/2007
WO	93/16798	9/1993	WO	2007/125332	A1	11/2007
WO	85/04867	11/1995	WO	2007/130054	A1	11/2007
WO	96/22263	7/1996	WO	2007/130055	A1	11/2007
WO	97/44302	11/1997	WO	2007/141295	A1	12/2007
WO	98/12165	3/1998	WO	2007/142745	A1	12/2007
WO	99/07443	2/1999	WO	2008/036562		3/2008
WO	00/07718	A1 2/2000	WO	2008/036563		3/2008
WO	00/09261	A1 2/2000	WO	2008/106319		9/2008

WO	2008/157043	12/2008
WO	2008/157044	12/2008
WO	2008/157045	12/2008
WO	2008/157046	12/2008
WO	2008/157047	12/2008

OTHER PUBLICATIONS

Abstract of JP2007099729, Method for producing alpha-methylstyrene or cumene, Publication date: Apr. 19, 2007, Inventor: Toshio, esp@cenet database—worldwide.

Abstract of RO119778, Process for preparing perchloroethylene, Publication date: Mar. 30, 2005, Inventor: Horia et al., esp@cenet database—worldwide.

Abstract of WO0105737, Method for preparing a carboxylic acid, Publication date: Jan. 25, 2001, Inventor: Pascal et al., esp@cenet database—worldwide.

Abstract of WO0105738, Method for preparing a carboxylic acid, Publication date: Jan. 25, 2001, Inventor: Pascal et al., esp@cenet database—worldwide.

Abstract of WO2004092099, Method for producing cyclic enols, Publication date: Oct. 28, 2004, Inventor: Marko et al., esp@cenet database—worldwide.

Abstract of WO2006063852, Electroluminescent polymers and use thereof, Publication date: Jun. 22, 2006, Inventor: Arne et al., esp@cenet database—worldwide.

Abstract of WO2006136135, Method for decarboxylating C-C cross-linking of carboxylic acids with carbon electrophiles, Publication date: Dec. 28, 2006, Inventor: Goossen Lukas et al., esp@cenet database—worldwide.

Abstract of WO2007028761, Method for chlorinating alcohols, Publication date: Mar. 15, 2007, Inventor: Rohde et al., esp@cenet database—worldwide.

Abstract of WO2007128842, Catalytic transalkylation of dialkyl benzenes, Publication date: Nov. 15, 2007, Inventor: Goncalvesalmeida et al., esp@cenet database—worldwide.

Abstract of WO2007137566, Method for catalytic conversion of organic oxygenated compounds from biomaterials, Publication date: Dec. 6, 2007, Inventor: Reschetilowski, esp@cenet database—worldwide.

Abstract of WO9721656, Method for making fluoroalknols, Publication date: Jun. 19, 1997, Inventor: Gillet, esp@cenet database—worldwide.

Abstract of WO9950213, Method for producing dialkyl ethers, Publication date: Oct. 7, 1999, Inventor: Falkowski Juergen et al., esp@cenet database—worldwide.

Adachi, et al.; Synthesis of Sialyl Lewis X Ganglioside Analogs Containing a Variable Length Spacer Between the Sugar and Lipophilic Moieties; *J. Carbohydrate Chem.*, vol. 17, No. 4-5, (1998), pp. 595-607, XP009081720.

Abstract of EP0039471, Process for the preparation of 2-chloro-1,1,1,2,3,3,3-heptafluoropropane, Publication date: Nov. 11, 1981, Inventor: Von Halasz, esp@cenet database—worldwide.

Abstract of EP0101337, Process for the production of methylene chloride, Publication date: Feb. 22, 1984, Inventor: Olah et al., esp@cenet database—worldwide.

Abstract of EP0407989, Method for the production of 1,1,1-trifluoro-2,2-dichloroethane by photochlorination, Publication date: Jan. 16, 1991, Inventor: Cremer et al., esp@cenet database—worldwide.

Abstract of EP0442258, Process for the preparation of a polyunsaturated olefin, Publication date: Aug. 21, 1991, Inventor: Gaudin et al., esp@cenet database—worldwide.

Abstract of EP0465294, Process for the preparation of unsaturated bromides, Publication date: Jan. 8, 1992, Inventor: Decaudin et al., esp@cenet database—worldwide.

Abstract of EP0549387, Synthesis of n-perfluorooctylbromide, Publication date: Jun. 30, 1993, Inventor: Drivon et al., esp@cenet database—worldwide.

Abstract of EP0850906, Process and apparatus for the etherification of olefinic hydrocarbon feedstocks, Publication date: Jul. 1, 1998, Inventor Masson, esp@cenet database—worldwide.

Abstract of EP0858987, Process for the conversion of lighter alkanes to higher hydrocarbons, Publication date: Aug. 19, 1998, Inventor: Amariglio et al., esp@cenet database—worldwide.

Bakker, et al.; An Exploratory Study of the Addition Reaction of Ethyleneglycol, 2-Chloroethanol and 1, 3-Dichloro-2-Propanol to 1-Dodecene; *J. Am. Oil Chem. Soc.*, vol. 44, No. 9 (1967), pp. 517-521; XP009081570.

Abstract of EP0235110, Process for the stabilization of silicalite catalysts, Publication date: Sep. 2, 1987, Inventor: Debras et al., esp@cenet database—worldwide.

Bouzide et al.; Highly Selective Silver (I) Oxide Mediated Monoprotection of Symmetrical Diols; *Tetrahedron Letters*, Elsevier, vol. 38, No. 34 (1997), pp. 5945-5948; XP004094157.

Combined International Search Report and Written Opinion Dated Apr. 17, 2007 for PCT/US06/13394, in the name of GRI, Inc.

Gibson; Phase-Transfer Synthesis of Monoalkyl Ethers of Oligoethylene Glycols; *Journal of Organic Chemistry*, vol. 45, No. 6 (1980) pp. 1095-1098; XP002427776.

Klabunde, Kenneth J., et al., Changes in Texture and Catalytic Activity of Nanocrystalline MgO during Its Transformation to MgC12 in the Reaction with 1-Chlorobutane, *J. Phys. Chem. B* 2001, 105, 3937-3941.

Loiseau et al.; Multigram Synthesis of Well-Defined Extended Bifunctional Polyethylene Glycol (PEG) Chains; *J. of Organic Chem.*, vol. 69, No. 3 (2004), pp. 639-647; XP002345040.

Mihai et al.; Application of Bronsted-type LFER in the study of phospholipase C mechanism; *J. Am. Chem. Soc.*, vol. 125, No. 11 (2003) pp. 3236-3242; XP002427799.

Motupally et al., Recycling Chlorine from Hydrogen Chloride: A New and Economical Electrolytic Process, *The Electrochemical Society Interface*, Fall 1998.

Nishikawa et al.; Ultrasonic Relaxations in Aqueous Solutions of Alcohols and the Balance Between Hydrophobicity and Hydrophilicity of the Solutes; *J. Phys. Chem.* vol. 97, No. 14 (1993), pp. 3539-3544; XP002427775.

Prelog et al.; Chirale 2,2'-Polyoxaalkano-9,9'-Spirobifluorene; *Helvetica Chimica ACTA*, vol. 62, No. 7, (1979) pp. 2285-2302; XP002030901.

Shimizu et al., Gas-Phase Electrolysis of Hydrobromic Acid Using PTFE-Bonded Carbon Electrode, *Int. J. Hydrogen Energy*, vol. 13, No. 6, pp. 345-349, 1988.

Velzen et al., HBr Electrolysis in the Ispra Mark 13A Flue Gas Desulfurization Process: Electrolysis in a DEM Cell, *J. of Applied Electrochemistry*, vol. 20, pp. 60-68, 1990.

Whitesides et al.; Nuclear Magnetic Resonance Spectroscopy. The Effect of Structure on Magnetic Nonequivalence Due to Molecular Asymmetry; *J. Am. Chem. Soc.*, vol. 86, No. 13 (1964), pp. 2628-2634; XP002427774.

JLM Technology Ltd.; "The Miller GLS Technology for Conversion of Light Hydrocarbons to Alcohols" *New Science for the Benefit of Humanity*; May 31, 2000; pp. 1-10.

Jaumain, Denis and Su, Bao-Lian; "Direct Catalytic Conversion of Chloromethane to Higher Hydrocarbons Over Various Protonic and Cationic Zeolite Catalysts as Studied by in-situ FTIR and Catalytic Testing"; 2000; pp. 1607-1612; *Studies in Surface Science and Catalysis 130*; Elsevier Science B.V.

Taylor, Charles E.; "Conversion of Substituted Methanes Over ZSM-Catalysts"; 2000; pp. 3633-3638; *Studies in Surface Science and Catalysis 130*; Elsevier Science B.V.

ZSM-5 Catalyst; <http://chemelab.ucsd.edu/methanol/memos/ZSM-5.html>; Nov. 6, 2003; p. 1.

Final Report; "Abstract"; <http://chemelab.ucsd.edu/methanol/memos/final.html>; May 9, 2004; pp. 1-7.

Driscoll, Daniel J.; "Direct Methane Conversion"; *Federal Energy Technology Center, U.S. Department of Energy*; M970779; 2001; pp. 1-10.

Olah et al.; "Selective Monohalogenation of Methane over Supported Acid or Platinum Metal Catalysts and Hydrolysis of Methyl Halides . . ."; *J. American Chemical Society* 1985, vol. 107; 0002-7863/85/1507-7097\$01.50/0; pp. 7097-7105.

Murray et al.; "Conversion of Methyl Halides to Hydrocarbons on Basic Zeolites: A Discovery by in Situ NMR"; *J. American Chemical Society* 1993, vol. 115; pp. 4732-4741.

- Lorkovic et al.; "A Novel Integrated Process for the Functionalization of Methane and Ethane: Bromine as Mediator", *Catalysis Today* 98; 2004; pp. 317-322.
- Lorkovic et al.; "C1 Oxidative Coupling via Bromine Activation and Tandem Catalytic Condensation over CaO/Zeolite Composites II . . ."; *Catalysis Today* 98; 2004; pp. 589-594.
- Olah et al.; "Antimony Pentafluoride/Graphite Catalyzed Oxidative Conversion of Methyl Halides with Copper Oxides (or Copper/Oxygen) to Dimethyl Ether"; *J. Org. Chem.* 1990, 55; 1990 American Chemical Society; pp. 4289-4293.
- Taylor, Charles E. et al.; "Direct Conversion of Methane to Liquid Hydrocarbons Through Chlorocarbon Intermediates"; 1988 Elsevier Science Publishers B.V. Amsterdam, Netherlands; pp. 483-489.
- Chang, Clarence D. et al.; "The Conversion of Methanol and Other O-Compounds to Hydrocarbons over Zeolite Catalysts"; *Journal of Catalysis* 47; 1977; Academic Press, Inc.; pp. 249-259.
- Zhou, Xiao-Ping et al.; "An Integrated Process for Partial Oxidation of Alkanes"; *Chem. Commun.* 2003; The Royal Society of Chemistry 2003; pp. 2294-2295.
- Sun, Shouli et al.; "A General Integrated Process for Synthesizing Olefin Oxides"; *Chem. Commun.* 2004; The Royal Society of Chemistry 2004; pp. 2100-2101.
- Lorkovic, Ivan M. et al.; "C1 Oxidative Coupling via Bromine Activation and Tandem Catalytic Condensation and Neutralization over CaO/Zeolite Composites II . . ."; *Catalysis Today* 98; 2004; pp. 589-594.
- Yilmaz, Aysen et al.; "Bromine Mediated Partial Oxidation of Ethane over Nanostructured Zirconia Supported Metal Oxide/Bromide"; *Microporous and Mesoporous Materials*, 79; 2005; pp. 205-214.
- Taylor, Charles E.; "PETC's On-Site Natural Gas Conversion Efforts"; Preprints of the Fuel Division, 208th National Meeting of the American Chemical Society, 39 (4); 1994; pp. 1228-1232.
- Ione et al.; "Syntheses of Hydrocarbons from Compounds Containing One Carbon Atom Using Bifunctional Zeolite Catalysts"; *Solid Fuel Chemistry (Khimiya Tverdogo Topliva)*; 1982; pp. 29-43; vol. 16, No. 6; Allerton Press. Inc.
- Olah, George A. et al.; "Hydrocarbons Through Methane Derivatives"; *Hydrocarbon Chemistry*; 1995; pp. 89-90; John Wiley & Sons, Inc.
- Akhrem, Irena S. et al.; "Ionic Bromination of Ethane and Other Alkanes (Cycloalkanes) with Bromine Catalyzed by the Polyhalomethane-2AIBr3 Aprotic Organic Superacids Under Mild Conditions"; *Tetrahedron Letters*, vol. 36, No. 51, 1995; pp. 9365-9368; Pergamon; Great Britain.
- Smirnov, Vladimir V. et al.; "Selective Bromination of Alkanes and Arylalkanes with CBr4"; *Mendeleev Commun.* 2000; pp. 175-176.
- Olah, George A.; "Electrophilic Methane Conversion"; *Acc. Chem. Res.* 1987, 20; pp. 422-428; American Chemical Society, Loker Hydrocarbon Research Institute and Dept. of Chemistry; University of Southern California.
- Olah, George A. et al.; "Antimony Pentafluoride/Graphite Catalyzed Oxidative Carbonylation of Methyl Halides with Carbon Monoxide and Copper Oxides (or Copper/Oxygen) to Methyl Acetate"; *J. Org. Chem.* 1990, 55; pp. 4293-4297; Loker Hydrocarbon Research Institute and Dept. of Chemistry; University of Southern California.
- Bagno, Alessandro et al.; "Superacid-Catalyzed Carbonylation of Methane, Methyl Halides, Methyl Alcohol, and Dimethyl Ether to Methyl Acetate and Acetic Acid"; *J. Org. Chem.* 1990, 55; pp. 4284-4289; Loker Hydrocarbon Research Institute; University of Southern California.
- Olah, George A. et al.; "Onium Ylide Chemistry. 1. Bifunctional Acid-Base-Catalyzed Conversion of Heterosubstituted Methanes into Ethylene and Derived Hydrocarbons. The Onium Ylide Mechanism of the C1-C2 Conversion"; *J. Am. Chem. Soc.* 1984, 106; pp. 2143-2149.
- Mochida, Isao et al.; "The Catalytic Dehydrohalogenation of Haloethanes on Solid Acids and Bases"; *Bulletin of the Chemical Society of Japan*, vol. 44; 1971; pp. 3305-3310.
- Richards, Ryan et al.; "Nanocrystalline Ultra High Surface Area Magnesium Oxide as a Selective Base Catalyst"; *Scripta Materialia*, 44; 2001; pp. 1663-1666; Elsevier Science Ltd.
- Sun, Naijian et al.; "Nanocrystal Metal Oxide—Chlorine Adducts: Selective Catalysts for Chlorination of Alkanes"; *J. Am. Chem. Soc.* 1999, 121; pp. 5587-5588; American Chemical Society.
- Mishakov, Ilya V. et al.; "Nanocrystalline MgO as a Dehydrohalogenation Catalyst"; *Journal of Catalysis* 206; 2002; pp. 40-48; Elsevier Science, USA.
- Wagner, George W. et al.; "Reactions of VX, GD, and HD with Nanosize CaO: Autocatalytic Dehydrohalogenation of HD"; *J. Phys. Chem. B* 2000, 104; pp. 5118-5123; 2000 American Chemical Society.
- Fenelonov, Vladimir B. et al.; "Changes in Texture and Catalytic Activity of Nanocrystalline MgO during Its Transformation to MgCl2 in the Reaction with 1-Chlorobutane"; *J. Phys. Chem. B* 2001, 105; pp. 3937-3941; 2001 American Chemical Society.
- <http://webbook.nist.gov/>; "Welcome to the NIST Chemistry Web-Book"; 2005; U.S. Secretary of Commerce on Behalf of the United States of America.
- Claude, Marion C. et al.; "Monomethyl-Branching of Long n-Alkanes in the Range from Decane to Tetracosane on Pt/H-ZSM-22 Bifunctional Catalyst"; *Journal of Catalysis* 190; 2000; pp. 39-48.
- Thomas, J. M. et al.; "Synthesis and Characterization of a Catalytically Active Nickel-Silicoaluminophosphate Catalyst for the Conversion of Methanol to Ethene"; *Chem. Mater.*; 1991, 3; pp. 667-672; 1991 American Chemical Society.
- Thomas, John Meurig et al.; "Catalytically Active Centres in Porous Oxides: Design and Performance of Highly Selective New Catalysts"; *Chem. Commun.*; 2001; pp. 675-687.
- Lorkovic, Ivan et al.; "C1 Coupling via Bromine Activation and Tandem Catalytic Condensation and Neutralization over CaO/Zeolite Composites"; *Chem. Commun.*, 2004; pp. 566-567.
- Tamura, Masuhiko et al.; "The Reactions of Grignard Reagents with Transition Metal Halides: Coupling, Disproportionation, and Exchange with Olefins"; *Bulletin of the Chemical Society of Japan*, vol. 44; Nov. 1971; pp. 3063-3073.
- Weissmermel, Klaus et al.; "Industrial Organic Chemistry"; 3rd Edition 1997. pp. 160-162, and 208.
- Abstract of BE812868, Aromatic hydrocarbons prodn. from chlorinated hydrocarbons, Publication date: Sep. 27, 1974, esp@cenet database—worldwide.
- Abstract of BE814900, Volatile aromatic cpds. prodn., Publication date: Sep. 2, 1974, esp@cenet database—worldwide.
- Abstract of CN1199039, Pentanol and its production process, Publication date: Nov. 18, 1998, Inventor: Kailun, esp@cenet database—worldwide.
- Abstract of CN1210847, Process for producing low carbon alcohol by directly hydrating low carbon olefines, Publication date: Mar. 17, 1999, Inventor: Zhenguo et al., esp@cenet database—worldwide.
- Abstract of CN1321728, Method for preparing aromatic hydrocarbon and hydrogen gas by using low-pressure gas, Publication date: Nov. 14, 2001, Inventor: Jie et al., esp@cenet database—worldwide.
- Abstract of CN1451721, Process for non-catalytic combustion deoxidizing coal mine gas for producing methanol, Publication date: Oct. 29, 2003, Inventor: Pengwan et al., esp@cenet database—worldwide.
- Abstract of CN1623969, Method for preparing 1, 4-benzene dimethanol, Publication date: Jun. 8, 2005, Inventor: Jiarong et al., esp@cenet database—worldwide.
- Abstract of CN1657592, Method for converting oil to multiple energy fuel product, Publication date: Aug. 24, 2005, Inventor: Li, esp@cenet database—worldwide.
- Abstract of CN1687316, Method for producing biologic diesel oil from rosin, Publication date: Oct. 26, 2005, Inventor: Jianchun et al., esp@cenet database—worldwide.
- Abstract of CN1696248, Method for synthesizing biologic diesel oil based on ion liquid, Publication date: Nov. 16, 2005, Inventor: Sun, esp@cenet database—worldwide.
- Abstract of CN1699516, Process for preparing bio-diesel-oil by using microalgae fat, Publication date: Nov. 23, 2005, Inventor: Miao, esp@cenet database—worldwide.
- Abstract of CN1704392, Process for producing alkylbenzene, Publication date: Dec. 7, 2005, Inventor: Gao, esp@cenet database—worldwide.

- Abstract of CN1724612, Biological diesel oil catalyst and method of synthesizing biological diesel oil using sai catalyst, Publication date: Jan. 25, 2006, Inventor: Gu, esp@cenet database—worldwide.
- Abstract of CN1986737, Process of producing biodiesel oil with catering waste oil, Publication date: Jun. 27, 2007, Inventor: Chen, esp@cenet database—worldwide.
- Abstract of CN100999680, Esterification reaction tech. of preparing biodiesel by waste oil, Publication date: Jul. 18, 2007, Inventor: Weiming, esp@cenet database—worldwide.
- Abstract of CN101016229, Refining method for bromomeoamyl alcohol, Publication date: Aug. 15, 2007, Inventor: Tian, esp@cenet database—worldwide.
- Abstract of DE3209964, Process for the preparation of chlorinated hydrocarbons, Publication date: Nov. 11, 1982, Inventor: Pyke et al., esp@cenet database—worldwide.
- Abstract of DE3210196, Process for the preparation of a monochlorinated olefin, Publication date: Jan. 5, 1983, Inventor: Pyke et al., esp@cenet database—worldwide.
- Abstract of DE3226028, Process for the preparation of monochlorinated olefin, Publication date: Feb. 3, 1983, Inventor: Pyke et al., esp@cenet database—worldwide.
- Abstract of DE3334225, Process for the preparation of 1, 2-dichloroethane, Publication date: Apr. 4, 1985, Inventor: Hebgen et al., esp@cenet database—worldwide.
- Abstract of DE4232056, 2,5-Di:methyl-hexane-2, 5-di:ol continuous prodn. from tert. butanol—by oxidative dimerisation in two phase system with vigorous stirring, using aq. phase with specified density to facilitate phase sepn., Publication date: Mar. 31, 1994, Inventor: Gnann et al., esp@cenet database—worldwide.
- Abstract of DE4434823, Continuous prodn. of hydroxy-benzyl alkyl ether, Publication date: Apr. 4, 1996, Inventor Stein et al., esp@cenet database—worldwide.
- Abstract of FR2692259, Aromatisation of 2-4C hydrocarbons—using a fixed-mobile-catalytic bed process, Publication date: Dec. 17, 1993, Inventor: Alario et al., esp@cenet database—worldwide.
- Abstract of FR2880019, Manufacturing 1, 2-dichloroethane, comprises cracking core hydrocarbonated source, separating into fractions, sending into chlorination reaction chamber and oxychlorination reaction chamber and separating from chambers, Publication date: Jun. 30, 2006, Inventor: Strebelle et al., esp@cenet database—worldwide.
- Abstract of FR2883870, Formation of 1, 2-dichloroethane useful in manufacture of vinyl chloride involves subjecting mixture of cracking products obtained by cracking of hydrocarbon source, to a succession of aqueous quenching, alkaline washing, and oxidation steps, Publication date: Oct. 6, 2006, Inventor: Balthasart et al., esp@cenet database—worldwide.
- Abstract of FR2883871, Preparing 1, 2-dichloroethane comprises cracking hydrocarbon to form mixture, sending mixture into storage reservoir, supplying mixture into chlorination and/or oxychlorination reactor, and separating 1, 2-dichloroethane from reactor, Publication date: Oct. 6, 2006, Inventor: Balthasart et al., esp@cenet database—worldwide.
- Abstract of IT1255246, Process for the preparation of dinitrodiphenylmethanes, Publication date: Oct. 20, 1995, Applicant: Enichem Spa et al., esp@cenet database—worldwide.
- Abstract of IT1255358, Process for the synthesis of 1, 4-butanediol, Publication date: Oct. 31, 1995, Inventor: Marco, esp@cenet database—worldwide.
- Abstract of JP2142740, Production of fluoroalcohol, Publication date: May 31, 1990, Inventor: Tsutomu et al., esp@cenet database—worldwide.
- Abstract of JP2144150, Chemical process and catalyst used therefore, Publication date: Jun. 1, 1990, Inventor: Deidamusu et al., esp@cenet database—worldwide.
- Abstract of JP4305542, Production of halogenated hydrocarbon compounds, Publication date: Oct. 28, 1992, Inventor: Shinsuke et al., esp@cenet database—worldwide.
- Abstract of JP6172225, Method for fluorinating halogenated hydrocarbon, Publication date: Jun. 21, 1994, Inventor: Takashi et al., esp@cenet database—worldwide.
- Abstract of JP6206834, Production of tetrachloroethanes, Publication date: Jul. 26, 1994, Inventor: Toshiro et al., esp@cenet database—worldwide.
- Abstract of JP8266888, Method for decomposing aromatic halogen compound, Publication date: Oct. 15, 1996, Inventor: Yuuji et al., esp@cenet database—worldwide.
- Abstract of JP2001031605, Production of 3-hydroxy-1-cycloalkene, Publication date: Feb. 6, 2001, Inventor: Hideo et al, esp@cenet database—worldwide.
- Abstract of JP2004075683, Method for producing optically active halogenohydroxypropyl compound and glycidyl compound, Publication date: Mar. 11, 2004, Inventor: Keisuke et al., esp@cenet database—worldwide.
- Abstract of JP2004189655, Method for fluorinating with microwave, Publication date: Jul. 8, 2004, Inventor: Masaharu et al., esp@cenet database—worldwide.
- Abstract of JP2005075798, Method for Producing adamantyl ester compound, Publication date: Mar. 24, 2005, Inventor: Norihiro et al., esp@cenet databas—worldwide.
- Abstract of JP2005082563, Method for producing 1, 3-adamantanediol, Publication date: Mar. 31, 2005, Inventor: Norihiro et al., esp@cenet database—worldwide.
- Abstract of JP2005145977, Process for catalytically oxidizing olefin and cycloolefin for the purpose of forming enol, olefin ketone, and epoxide, Publication date: Jun. 9, 2005, Inventor: Cancheng et al., esp@cenet database—worldwide.
- Abstract of JP2005254092, Method of manufacturing alkynes, Publication date: Sep. 22, 2005, Inventor: Shirakawa Eiji, esp@cenet database—worldwide.
- Abstract of JP2006151892, Preparation method of alcohol derivative, Publication date: Jun. 15, 2006, Inventor: Baba Akio et al., esp@cenet database—worldwide.
- Abstract of JP2006152263, Organic-inorganic hybrid-type mesoporous material, method for producing the same, and solid catalyst, Publication date: Jun. 15, 2006, Inventor: Junko et al., esp@cenet database—worldwide.
- Abstract of JP2006193473, Aryl polyadamantane derivative having carboxy or acid anhydride group and method for producing the same, Publication date: Jul. 27, 2006, Inventor: Yasuto et al., esp@cenet database—worldwide.
- Abstract of JP2006231318, Phosphorus containing macromolecule immobilizing palladium catalyst and method for using the same, Publication date: Sep. 7, 2006, Inventor: Osamu et al., esp@cenet database—worldwide.
- Abstract of JP2006263567, Optical resolution method of optical isomer and optical resolution device, Publication date: Oct. 5, 2006, Inventor: Yoshikazu et al., esp@cenet database—worldwide.
- Abstract of JP2006265157, Method for catalytically activating silicated nucleating agent using phosphazene base, Publication date: Oct. 5, 2006, Inventor: Yoshinori et al., esp@cenet database—worldwide.
- Abstract of JP2006306758, Method for producing biaryl compound, Publication date: Nov. 9, 2006, Inventor: Yuji et al., esp@cenet database—worldwide.
- Abstract of JP2007001942, Production method of para-xylene, Publication date: Jan. 11, 2007, Inventor: Kazuyoshi, esp@cenet database—worldwide.
- Abstract of JP2007015994, Method for synthesizing organic compound in ultra high rate under high temperature and high pressure water, and system of high temperature and high pressure reaction, Publication date: Jan. 25, 2007, Inventor: Hajime et al., esp@cenet database—worldwide.
- U.S. Office Action from U.S. Appl. No. 10/826,885 dated Oct. 31, 2005.
- U.S. Office Action from U.S. Appl. No. 10/826,885 dated Apr. 19, 2006.
- U.S. Office Action from U.S. Appl. No. 10/826,885 dated Jul. 27, 2006.
- U.S. Office Action from U.S. Appl. No. 10/826,885 dated Nov. 2, 2006.
- U.S. Office Action from U.S. Appl. No. 10/826,885 dated Jan. 24, 2007.

US 8,415,512 B2

Page 10

U.S. Office Action from U.S. Appl. No. 11/101,886 dated Jan. 24, 2007.

U.S. Office Action from U.S. Appl. No. 11/254,438 dated Jan. 24, 2007.

U.S. Office Action from U.S. Appl. No. 11/254,438 dated Nov. 1, 2007.

U.S. Office Action from U.S. Appl. No. 10/893,418 dated Jan. 2, 2008.

U.S. Office Action from U.S. Appl. No. 10/893,418 dated Jun. 14, 2007.

U.S. Office Action from U.S. Appl. No. 11/091,130 dated Oct. 3, 2007.

U.S. Office Action from U.S. Appl. No. 10/365,346 dated Jun. 12, 2006.

U.S. Office Action from U.S. Appl. No. 11/103,326 dated Aug. 31, 2007.

U.S. Office Action from U.S. Appl. No. 11/103,326 dated Dec. 6, 2006.

U.S. Office Action from U.S. Appl. No. 11/098,997 dated Nov. 20, 2008.

U.S. Office Action from U.S. Appl. No. 12/215,326 dated Feb. 10, 2009.

U.S. Office Action from U.S. Appl. No. 10/430,240 dated Mar. 6, 2006.

U.S. Office Action from U.S. Appl. No. 10/369,148 dated Oct. 16, 2006.

U.S. Office Action from U.S. Appl. No. 10/369,148 dated Mar. 14, 2006.

U.S. Office Action from U.S. Appl. No. 10/894,165 dated Aug. 16, 2006.

U.S. Office Action from U.S. Appl. No. 12/080,594 dated Dec. 22, 2008.

U.S. Office Action from U.S. Appl. No. 11/703,358 dated Jun. 11, 2008.

* cited by examiner

HYDROCARBON CONVERSION PROCESS IMPROVEMENTS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 12/692,831 filed Jan. 25, 2010 now U.S. Pat. No. 7,838,708, which is a continuation of U.S. patent application Ser. No. 11/098,997 filed Apr. 5, 2005 now abandoned, which is a continuation-in-part of U.S. patent application Ser. No. 10/430,240 filed Aug. 19, 2003 now U.S. Pat. No. 7,161,050, which is a continuation-in-part of prior application Ser. No. 10/365,346 filed Feb. 12, 2003 now abandoned, which is a continuation of prior application Ser. No. 10/298,440 filed Nov. 19, 2002, abandoned, which is a continuation-in-part of prior application Ser. No. 10/208,068, filed Jul. 29, 2002, abandoned, which is a continuation-in-part of prior application Ser. No. 10/054,004 filed Jan. 24, 2002, now U.S. Pat. No. 6,486,368, which is a continuation-in-part of prior application Ser. No. 09/951,739, filed Sep. 11, 2001, now U.S. Pat. No. 6,465,696, which is a continuation-in-part of application Ser. No. 09/886,078 filed Jun. 20, 2001, now U.S. Pat. No. 6,472,572.

CLAIM OF PRIORITY

Applicant claims priority based on provisional patent application Ser. No. 60/559,844, filed Apr. 6, 2004.

TECHNICAL FIELD

This invention relates generally methods and apparatuses for synthesizing olefins, alcohols, ethers, and aldehydes from alkanes, alkenes, and aromatics, and more particularly to specific improvements in the methods and apparatuses disclosed in the patents and patent applications identified herein.

BACKGROUND AND SUMMARY OF THE INVENTION

The present application comprises a continuation-in-part of application Ser. No. 10/430,240, the disclosure of which is incorporated herewith by reference as if set forth herein. The present invention comprises specific improvements in and to the methods and apparatuses disclosed and described in the patents and patent applications identified herein, specifically including:

- Safety Improvements;
- Use of Alternative Feedstocks;
- Process Simplification;
- Improvements to the Halogenation Step;
- Improvements to the Reproportionation Step;
- Improvements to the Solid Oxide Reaction;
- Improvements to Solid Oxide Regeneration;
- Improvements in Separations;
- Maintenance;
- Start-up;
- Shut-down;
- Materials of Construction.

DETAILED DESCRIPTION

Safety Improvements:

1. The safety of the process may be improved by shipping the halogen in solid halide form. Some specific variations include:

a. The solid may be one of the solids that will be used in the process. One advantage is that separate halide liberation

equipment may be avoided. The solid may be regenerated using air, oxygen, and/or oxygen-containing gas in the process equipment.

- b. The solid may also be a solid that is not used in the process, but rather a solid which is inexpensive, relatively non-toxic, able to liberate halide at lower temperature, liberates halide with heating rather than oxidation, less hygroscopic, less corrosive in solid form, less corrosive to regenerate, less volatile, more dense, containing a higher halogen content, disposable, and/or otherwise more convenient for shipping. Examples include:
- i. Copper chloride, copper bromide, or copper iodide may be used for the shipment of chlorine, bromine, or iodine due to relatively low cost of copper relative to some other materials.
 - ii. Copper halides may be used for shipment for processes in which the reactive solid contains nickel, chromium, lead, cobalt, or other potentially toxic elements.
 - iii. Calcium bromide may be used to ship bromine for use in a process using a cobalt-containing metal oxide since cobalt bromide readily forms highly hydrated species.
 - iv. Copper chloride, copper bromide, or copper iodide may be used for the shipment of chlorine, bromine, or iodine since copper halides may be easily regenerated with air, oxygen, or oxygen containing gas at temperatures below those required to regenerate other solids.
 - v. Copper (II) bromide may be used to ship bromine since bromine may be liberated by heating without oxygen.
 - vi. Copper bromide may be used to ship halide for processes using iron-containing materials since iron bromide is hygroscopic, potentially volatile, and potentially corrosive.
 - vii. Pure copper halide may be used to ship halide for processes using a supported metal oxide. Such a choice will eliminate the need to transport the inert support.
 - viii. Carbon tetrabromide may be used with combustion of this material either in dedicated equipment or within a process unit, generating bromine and carbon dioxide. Such a solid is disposable, thereby eliminating the requirement of returning the solid oxide to a processing facility.
- c. The solid may also be a solid that is not used in the process so that the solid used may be shipped in oxide form, which may be more stable, less likely to sinter, dissolve, adsorb (or absorb) water (or other contaminants), or fracture.
- d. As the solid used in the process can hold substantially more bromine than the optimum level for use in the process, the solid could be shipped to an operating facility with a relatively high level of bromine that could be used to make up the small amount of bromine that may be lost during normal operation of the process.
2. The safety of the process may be improved by shipping the halogen in liquid halide form. The use of liquid may minimize solids handling operation and associated particulate hazards. Liquid may also be easier to handle.
- a. Specifically, boron tribromide may be used with oxidation to boron oxide liberating bromine.
3. The safety of the process may be improved by shipping the halogen in liquid alkyl halide form. The use of liquid halides may minimize solids handling operation and associated particulate hazards. The use of liquid halides may also be

3

easier to handle. The shipment of alkyl halide may be particularly useful for the startup of the plant and may also provide a convenient and safe way to introduce make-up halide.

4. The safety of the process may be improved by the placement of hygroscopic metal halides in selected reactors provided with a sink for water in the event of a process upset. Many metal halides are hygroscopic and will react with water to form hydrates, minimizing corrosion.

a. The metal halides may be selected and placed so that they are molten in the hydrated form and easily removed.

b. The metal halides may be selected based on a low melting temperature so that they can be pumped into the process in the event of an upset.

5. The bromine inventory may be reduced by utilization of the bromine separation apparatus (typically following the regeneration reactor) as the reservoir for bromine for introduction into the alkane bromination reactor or other necessary step involving bromine. This reservoir of liquid bromine will have sufficient capacity to maintain adequate pump priming and allow bromine to be pumped as a liquid rather than using more costly compressors.

6. Reactive metal oxide traps at all process vents for use in normal and emergency operations may be used to insure against release of any and all organic-bromides. These metal oxides may be regenerated to recover bromine.

7. The safety of the process may be enhanced by the use of a solid oxide to dispose of halogenated organic streams and recovery of halide by conversion to carbon dioxide, water, and solid halide. The solid oxide may be regenerated by reacting the solid halide with oxygen, liberating halogen for recycle to the process.

a. One example is the use of CuO or CuZrO₃ to convert vinyl bromide to carbon dioxide, water, coke, and CuBr or CuBrZrO₂.

Use of Alternate Feedstocks

The above-identified processes may be useful and particularly valuable with feedstocks containing otherwise difficult to separate components. The halogenation chemistry may facilitate the reactive separation of various streams including:

1. The use of streams containing alkane and olefin of the same carbon number.

a. The olefin may be converted with molecular halogen or solid halide to the 1,2-dihalide for use as a feedstock to an epoxide process. The 1,2-dihalide will be easy to separate from the alkane. Examples include:

i. Converting the propylene in a stream containing propane and propylene to 1,2-dibromopropane and subsequently to propylene oxide. The 1,2-dibromopropane is formed by reacting the mixed hydrocarbon stream with bromine, most preferably at low temperatures where little appreciable reaction with propane occurs. Separation of propane from propylene is required in many existing plants including ethylene plants and is considered one of the most difficult separations in the chemical industry.

b. The olefin may be converted with wet halogen to halohydrin for use as a feedstock to an epoxide process. The halohydrin will be easy to separate from the alkane. Examples include:

i. Converting the propylene in a stream containing propane and propylene to the bromohydrin and subsequently to propylene oxide. The propylene is converted by passing the hydrocarbon stream through bromine water.

c. The olefin may be converted with hydrogen halide to form the monohalide for use as a feedstock in an olefin,

4

alcohol, epoxide, aldehyde, ketone, or other process. The halide will be easy to separate from the alkane. Examples include:

i. Converting olefin in a gasoline feed to alkyl halide by reacting with hydrogen halide to form alkyl halide. The alkyl halide can be easily removed, leaving olefin-depleted gasoline.

ii. Converting butenes to butyl halides in a mixed feed of butanes and butenes. The butyl halides may be coupled to products containing eight carbon atoms for use in gasoline.

2. The use of streams containing branched and linear alkanes resulting in product streams enriched in branched and/or linear molecules.

a. Branched alkanes containing tertiary carbon may be selectively halogenated to alkyl halide and separated, leaving a stream enriched in linear alkane. Examples include:

i. Depletion of the branched content of detergent-range alkanes by reaction with halide, resulting in greater reactivity with the branched alkanes to branched alkyl halides. Following separation of the branched halides, the remaining stream is enriched in linear alkane. The stream rich in branched halides may be dehydrohalogenated either catalytically or using a solid oxide to create a stream rich in branched olefin for hydroformylation and conversion to branched alcohols.

b. Branched alkanes containing tertiary carbon may be selectively halogenated to alkyl halide. Following separation, the halide stream will inevitably contain some non-branched halides. By selectively dehydrohalogenating the tertiary halides, a stream containing a very high fraction of branched olefins can be separated from the remaining halides. The selective dehydrohalogenation may be conducted thermally, using a catalyst at temperatures below those required for secondary alkyl halide dehydrohalogenation, or using a solid oxide catalyst at temperatures below those required for secondary alkyl halide dehydrohalogenation. Dehydrohalogenating the remaining halides will leave a stream enriched in linear olefins.

3. The use of streams containing multiple types of branched molecules resulting in product streams enriched or depleted in molecules containing a certain type or amount of branching:

a. Streams containing linear, mono-branched, and multiply-branched alkanes may be enriched or depleted in multiply-branched product by halogenating the multiply-branched alkanes to multiply-halogenated separation. The multiply-halogenated and/or mono-halide species may be easily separated. Following the desired separation of the non-halogenated, mono-halogenated, and multi-halogenated species, and dehydrohalogenation of the halides, the various streams may be recombined to generate the desired branching composition.

b. Streams containing branched alkanes with and without multiple branching at a single carbon (quaternary carbon) may be depleted in these quaternary carbon-containing species by halogenating the branched alkanes without the quaternary carbon, separating these halides, and dehydrohalogenating. The result will be streams rich in branched olefins without quaternary carbon and alkanes with quaternary carbon.

4. The use of streams containing trace amounts of impurities that are more reactive than the desired alkane reactant:

a. Alkane streams containing aromatics, alcohols, olefins, aldehydes, ketones, sulfides, sulfates, or other reactive

5

- molecules may be halogenated at low temperature to selectively halogenate the impurities for removal.
- b. Streams of mixed alkanes (e.g. natural gas, refinery streams) may be differentially halogenated based on differing rates of halogenation and subsequently reacted with metal oxides at lower temperatures where the non-halogenated alkanes would pass through without reaction.
5. The use of streams of mixed alkane and olefin in coupling processes. Streams containing alkanes and olefins may be used to produce products of higher carbon number. Several process variations may be employed:
- a. A process with:
 - i. Olefin hydrohalogenation in the presence of the alkane;
 - ii. Separation of the resulting alkyl halide from the alkane;
 - iii. Halogenation of Alkane;
 - iv. Separation of the resulting alkyl halide from the alkane;
 - v. Recycle of alkane;
 - vi. Feed of the alkyl halide to the coupling reactor.
 - b. Several variations of (a) may be employed:
 - i. Specifically, step ii (separation of alkyl halide after hydrohalogenation) may or may not be omitted.
 - ii. The alkane may or may not be separated from the alkyl halide (step iv).
 - iii. Reproportionation chemistry may or may not be employed.
 - iv. The olefin and alkane may be separated at the beginning of the process.
 - v. Halogenation may precede hydrohalogenation, particularly if high temperature is employed to hinder addition of halogen to the olefin.
 - vi. The hydrogen halide used for hydrohalogenation may or may not be the same formed in the halogenation step.

Process Simplification

1. The halogenation and solid oxide reaction steps may be conducted in the same unit:
 - a. The halogenation and solid oxide reaction may be simultaneous.
 - b. The halogenation may occur first by varying the contacting of the hydrocarbon, halogen, and solid oxide.
2. The halogenation, solid oxide reaction, and solid oxide regeneration may be conducted in the same unit by introducing hydrocarbon and oxygen to a solid halide or solid halide-oxide combination. The oxygen will regenerate the solid halide generating hydrocarbon halide and solid oxide, the hydrocarbon halide will react with the oxide, generating product. Variations include:
 - a. Periodic switching of the direction of feed to the reactor to minimize halogen migration from the reactor. Examples include:
 - i. Coupling methane to heavier products by cofeeding methane and oxygen over a metal-halide-containing solid.
3. The solid oxide reaction and product separation may be conducted simultaneously when the product is lighter than the reactant the reaction is conducted in a liquid phase reactor under conditions where the product is a vapor and leaves the reactant mixture.
4. Reacting the halide-containing regeneration effluent with olefin to form dihaloalkanes to reduce the energy required for and equipment size in the halide recovery.
5. Reacting the alkane over selected metal-halides in the regeneration step to form the alkyl-halide and a metal-hy-

6

dride. This would also be a safety improvement and eliminate the need for halogen separation. Materials include but are not limited to halides of boron, nickel, iron, and their mixtures as well as carbon based materials (e.g. C60).

6. Operation of the halogenation process at high halogen: alkane ratio for the feed at temperatures and pressures to maximize the production of monohalo-alkanes at 100% alkane conversion. The alkane feed may be mixed. The products which will contain multiply-halogenated species and haloacid which may be passed directly over a metal oxide bed to produce a mixture of products dependent upon the reaction conditions will be produced which will be condensed together and separated in the liquid phase by a combination of distillation and phase separation.

7. The use of a hydrogenation step to recover the over-halogenated products by reducing the halogenation to the desired degree. The use of such a step will allow for higher per-pass conversion in the halogenation step. Catalysts may be used, including but not limited to Pd, Pt, Ru, Ni, Au, Cu, and their alloys.

8. Controlling the amount of hydrogen halide added to a metal oxide reactor in order to generate the heat required for an endothermic reaction.

9. The use of hydrogen halide formed in the halogenation step for conversion of byproducts or products into more useful compounds.

- a. For example, HBr could be used to hydrobrominate vinylbromide, a common undesirable by-product resulting from HBr elimination from dibromoethane back to same (or any higher vinylbromide equivalent to the corresponding dibromoalkane). In another example, HBr could be used in the acidic cleavage of ethers into alcohols and alkylbromides, the former increasing the yield of the desired alcohol product and the latter being recycled to the educt stream for reaction on the metal oxide.

Improvements to the Halogenation Step

1. Improvements in selectivity to desired multiply-halogenated isomers through isomerization of the multiply-halogenated species formed by halogenation. Examples include:
 - a. Forming dihalides dehydrohalogenating the dihalides, and rehydrohalogenating to form the desired isomers. The rehydrohalogenation may be conducted using process conditions different from the initial halogenation to enhance yield of the desired isomer. The process conditions varied may include temperature, pressure, and catalyst. Some examples include:
 - i. Halogenating ethane so that it contains mixed halides including 1,1 and 1,2-dihaloethanes. Dehydrohalogenating the dihaloethanes, and rehydrohalogenating to enrich the 1,2-dihaloethane content.
 - ii. Halogenating propane so that it contains mixed halides including 1,1, 2,2, 1,3, and 1,2-dihalopropanes. Dehydrohalogenating the dihalopropanes, and rehydrohalogenating to enrich the 1,2-, 2,2-, 1,3, or 1,1-isomer content.
 - iii. Halogenating butane so that it contains mixed dihalides. Dehydrohalogenating, and rehydrohalogenating to enrich the 2,2- or 2,3-isomer content. The 2,2- or 2,3-isomer may be reacted with a metal oxide to make methyl-ethyl ketone.
 - iv. Halogenating butane so that it contains mixed tetrahalides. Dehydrohalogenating, and rehydrohalogenating to enrich the 1,2,3,4-isomer content.
 - v. Halogenating cyclohexane so that it contains mixed halides including 1,1, 1,2, 1,3, and 1,4-dihaloisomers.

Dehydrohalogenating, and rehydrohalogenating to enrich the 1,1, 1,2, 1,3, and 1,4-dihaloisomer content.

2. Enrichment in the primary halide content of a stream of mixed halide isomers by separating primary halides from other halide isomers. Dehydrohalogenating the other isomers, rehydrohalogenating the resulting olefins to produce a stream enriched in primary halide isomers, and returning the resulting stream to the primary halide separation step. Some variations include:

- a. Using selective dehydrohalogenation of the non-primary isomers to form easily separated olefin and hydrogen halide.
- b. Using distillation to separate the primary and other isomers.
- c. Using adsorption to separate the primary and other isomers.
- d. Using a shape-selective catalyst to rehydrohalogenate the olefin, enhancing primary halide yield.

3. The use of multiple halogens to create the desired halide isomer. One halogen may be used to halogenate the hydrocarbon and be replaced by another.

4. The use of a membrane reactor with halogen on one side and alkane on the other to improve selectivity to the desired halide isomer. This reactor design may improve monohalogenation, dihalogenation, and/or primary halogenation selectivity.

5. Operation of the halogenation reaction at high halogen:alkane ratio to improve conversion may result in unconverted halogen. Photoactivation of the unconverted halogen may be used at low temperature in a solid oxide bed to allow full recovery of all the halogen.

Improvements to the Reproportionation Step

In many processes, the overhalogenated species may be recycled to a point in the process where they are converted to the desired degree of halogenation or less than the desired degree of halogenation. The change in degree of halogenation is termed "reproportionation," and allows for the use of the carbon and hydrogen in the overhalogenated species, thus reducing feedstock loss and perhaps also allowing greater economic per-pass yield.

Several Improvements Include:

1. A low-temperature reproportionation step, in which the halogen is redistributed among over-halogenated species, resulting in the formation of optimally halogenated species and additional very highly halogenated species.

a. An example is a mixture of dibromomethane, tribromomethane, and tetrabromomethane are allowed to react, producing a stream enriched in methyl bromide and tetrabromomethane.

2. A low-temperature reproportionation step, in which the halogen is redistributed among over-halogenated species, resulting in the formation of optimally halogenated species and additional very highly halogenated species. The yield of optimally halogenated species is maximized by conducting this reproportionation under temperature, pressure, and process conditions such that the reproportionation is conducted in the liquid phase while the optimally halogenated species is predominantly in the vapor phase.

a. An example is: a mixture of liquid dibromomethane, tribromomethane, and tetrabromomethane are allowed to react in the presence of a catalyst at about 30 C. As the bromine is redistributed and methyl bromide is formed, much of the methyl bromide leaves the solution and enters the vapor phase.

3. The conversion of over-halogenated hydrocarbon to carbon black or other carbon material and halogen. The carbon material may be sold and the halogen may be recycled to the process.

4. The reproportionation of overhalogenated hydrocarbon with another hydrocarbon or haloalkane. Such a process may allow the recovery of the desired hydrocarbon with a loss of a less desirable material.

a. For example, dibromomethane is reacted with propane to make methyl bromide and brominated propanes. Ideally, one propane molecule can be used to convert eight dibromomethane molecules to methyl bromide. The bromine can be recovered from the brominated propane through thermal decomposition, oxidation, reaction with solid oxide, or other means.

Improvements to the Solid Oxide Reaction

1. A method of contacting water with alkyl halide and metal oxide in a multi-phase reactor with alkyl halide, solid oxide and optional diluent present at the bottom of the reactor with refluxing water present in a zone above the reactant mixture.

2. The product yield may be increased and process corrosivity may be reduced by conducting the solid oxide reaction in a liquid phase with water present to remove metal halide as it is formed. A specific example is:

a. The reaction is conducted in a vessel containing liquid alkyl halide, liquid water, water vapor, and solid. The water vapor condenses at the top of the reactor or is returned from an external condenser and settles through the metal oxide and alkyl halide containing phase. The water dissolves metal halide as it passes through the alkyl halide phase. The solid oxide may be supported on a plate to keep it out of the liquid water phase. The water and metal halide passes into a separate liquid phase at the bottom of the reactor where some of the water is vaporized. Variations include:

i. A batch reactor.

ii. A continuous reactor in which alkyl halide, metal oxide, and water (or steam) are added continuously and metal halide solution is removed continuously to a regeneration reactor where it is dried and regenerated.

iii. The use of precipitation to remove metal halide from the metal halide solution. By reducing the temperature of the solution, some of the metal halide will precipitate for regeneration. The depleted metal halide solution may be recovered by filtration, centrifugation or other solids-liquid separation methods and recycled to the reactor. Recovered solids can be dried and regenerated to metal oxide and bromine.

3. The liquid phase performance of a reactor may be improved by adding a diluent. The diluent may be, but is not limited to alkanes that are readily separated from the products and reactants.

4. The yield to desired product may be improved by introducing the stream containing hydrocarbon halide to the metal oxide in stages.

5. The yield to desired product may be improved by providing a feed of solid to a fluidized bed reactor that includes some partially or completely spent material. Spent is defined as solid with no remaining oxygen (donation) capacity or bromine capacity.

6. The yield to desired product may be improved by providing a feed of solid to a fluidized bed reactor that includes some partially coked material.

7. The solid oxide reaction may be conducted in a series of switched fixed beds, some of which are undergoing regeneration at any given time.

8. In a process for the production of olefins, the di-halogenated species may be at least partially converted to olefin using certain solids. Some examples include:

- a. The reaction of silver metal with 1,2-dibromoethane to form ethylene and silver bromide. The silver bromide may be decomposed to silver and bromine using heat or electromagnetic radiation.
- b. The reaction of copper (I) bromide with 1,2-dibromomethane to form ethylene and copper(II) bromide. The copper (II) bromide may be decomposed to copper (I) bromide and bromine using heat.
- c. The reaction of 1,2-dibromomethane with a metal oxide to form ethylene, carbon dioxide, water, and metal bromide. The metal bromide may be regenerated by reaction with oxygen.

Improvements to Solid Oxide Regeneration

1. Varying the temperature of solid oxide prior to oxygen introduction to change the particle size of the solid oxide to a more desirable distribution.

- a. By raising the temperature, particularly to that above the regeneration onset temperature, prior to introduction of oxygen or air, the metal oxide obtained after regeneration may be reduced in the amount of fines or agglomerates it contains.
- b. By introducing oxygen at low temperature, particularly at that below the regeneration onset temperature, the metal oxide obtained after regeneration may be reduced in the amount of fines or agglomerates it contains.

2. Increasing the temperature of solid oxide prior to oxygen introduction to dehydrogenate or desorb adsorbed hydrocarbon, reducing the amount of water and possibly carbon oxides generated in regeneration, thus reducing corrosivity and simplifying halide purification.

3. Performing a separate oxidation, particularly at low temperature, to remove adsorbed hydrocarbon reducing the amount of water and carbon oxides generated in regeneration, thus reducing corrosivity and simplifying halide purification.

4. Introducing water to the solid halide to change the particle size of the resulting solid oxide to a more desirable distribution.

- a. The water may be introduced in the gas phase.
- b. The water may be introduced in the liquid phase.
- c. The water may be introduced concurrently with or prior to the introduction of oxygen.
- d. The hydrated solid may be allowed to settle and agglomerate.
- e. The hydrated solid may be subjected to intense fluidization to break apart agglomerates.
- f. A slurry or aqueous phase may be formed and dried in a manner to form the desired particle size. In particular, spray drying may be used.

5. Dissolving the active metal halide to separate it from impurities, and then converting metal halide to metal oxide.

6. The use of very high temperature regeneration to remove impurities. In particular, chlorine may be removed from metal bromide in this manner.

a. The combination of high temperature with heating of the solid halide prior to oxygen introduction may be particularly useful. In the case of metal bromides, this methodology may allow the removal of chlorine as CIBr or Cl₂.

7. The reduction of the solid halide with hydrogen or other reducing agent to remove impurities. The reduced material may be reoxidized with oxygen, air, or other oxygen containing gas.

Improvements in Separations

1. Separation of halogen from nitrogen, oxygen, and other non-condensibles using solid adsorbents. The solid adsor-

bents will adsorb the halogen, which can be removed by heating the solid or reducing the pressure. The adsorbents may be, but are not limited to:

- a. Molecular sieves;
- b. Mesoporous materials;
- c. Zeolites;
- d. Silica;
- e. Alumina;
- f. Aluminosilicates;
- g. Magnesia;
- h. Activated carbon;
- i. Metal bromides;
- j. Metal oxides;

2. Separation of halogen from nitrogen, oxygen, and other non-condensibles using reactive solid adsorbents. The solid reactive adsorbents will react with the halogen, forming a new chemical composition, from which the halogen can be removed by heating the solid or reducing the pressure, regenerating the solid. The reactive adsorbents may be, but are not limited to:

- a. Copper (I) bromide;
- b. Iron (II) bromide;
- c. Silver bromide;
- d. Carbon;
- e. Carbon, particularly fullerenes or nano-tubular carbon.

3. Removal of water from halogen by passing the mixed stream over metal halides or metal halide hydrates which may be supported or unsupported. The metal halides will form hydrates and the metal halide hydrates will form more highly hydrated species. The water can be liberated and starting material can be regenerated by heating.

4. Methods of removing trace amounts of halogen from product streams using reactive solids, which may or may not be regenerable. Some specific reactive solids include, but are not limited to:

- a. Copper (II) oxide;
- b. Silver;
- c. Copper;
- d. Lithium;
- e. Magnesium;
- f. Alkali metals.

5. Removal of residual halogen from streams by reaction with olefins. Specific examples include:

- a. Reacting with ethylene or propylene to form dihaloalkane, which can be converted to epoxide.
- b. Reacting with a heavy multiple-olefin to form highly halogenated species from which the halide can be recovered.

6. Removal of residual hydrocarbon halide from streams by reaction with reactive solids. Some specific reactive solids include, but are not limited to:

- a. Copper (II) oxide;
- b. Silver;
- c. Copper;
- d. Lithium;
- e. Magnesium;
- f. Alkali metals.

7. Separation of primary, secondary, and/or tertiary alkyl halides by selective dehydrohalogenation of selected species, separation of the olefin and hydrogen halide from the remaining alkyl halide, and recombination of the hydrogen halide and olefin to form alkyl halides. Examples include, but are not limited to:

- a. Separation of primary from secondary alkyl halides by selectively dehydrohalogenating the secondary alkyl halides to olefin and hydrogen halide, separation of the olefin and hydrogen halide from the primary halide, and

11

recombination of the hydrogen halide and olefin to form secondary and possibly primary halide.

- b. Separation of primary and secondary alkyl halides from tertiary alkyl halides by selectively dehydrohalogenating the tertiary alkyl halides to olefin and hydrogen halide, separation of the olefin and hydrogen halide from the primary and secondary alkyl halide, and recombination of the hydrogen halide and olefin to form alkyl halide.

8. Removal of sulfur-containing compounds from a hydrocarbon feed by reacting the feed with dry halogen to form sulfur, which can be removed as a solid from the sulfur-depleted hydrocarbon and hydrocarbon halide stream.

9. Removal of sulfur-containing compounds from a hydrocarbon feed by reacting the feed with dry halogen to form sulfur, which can be removed as a solid from the sulfur-depleted hydrocarbon and hydrocarbon halide stream.

10. Removal of carbon dioxide from a stream by reacting with a carbonate-forming material such as calcium oxide. The carbonate may be used in a hydrogen-halide recovery section of the plant. The carbonate will react with hydrogen halide, liberating water and carbon dioxide and producing solid halide, which can be regenerated and recycled to the carbon dioxide separation section.

11. Removal of arsenic, mercury, heavy metal-containing compounds from a hydrocarbon feed by reacting the feed with dry halogen to form solid metal compounds, solid metal halides or halogenated metal hydrocarbons, which can be easily separated.

12. Removing adsorbed product from the solid by rinsing with a compound that is easily separated from the product. Such a rinsing agent may be pentane or other alkane.

13. Removing adsorbed product from the solid by steam distillation.

Maintenance

1. A method of removing coke from reactors by reacting with bromine to form volatile carbon bromides. The carbon bromides may be used in the process in a reproporationation step, thus producing product from the coke.

2. A method of removing coke from reactors by reacting with hydrogen bromide to form volatile hydrocarbon bromides. The carbon bromides may be used in the process in a reproporationation step, thus producing product from the coke.

Start-Up

1. Starting the process with some or all of the solid in the halide or partially halogenated form may provide a number of benefits including:

- The ability to start the regeneration reactor early in the start up sequence.
- Reduction in the amount of oxygen carried over into the bromine separation unit.
- Reduction in the heat generated in the metal oxide reactor or hydrogen halide neutralization step.
- Reduction in the amount of adsorbed hydrocarbon and thus reduction in the amount of water and carbon dioxide generated in the regeneration unit.
- Reduction in unfavorable changes in the particle size distribution of the metal oxide.
- Improvement in the packing of a fixed bed reactor.
- Providing a hygroscopic metal halide to reactively remove water during upsets.

2. Starting the process with some or all of the solid in the oxide or partially oxygenated form may provide a number of benefits including:

- The ability to start the solid oxide reactor early in the start up sequence.

12

- Reduction in the amount of bromine generated in the regeneration reactor.

- Reduction in the heat generated in the metal oxide reactor or hydrogen halide neutralization step.

- Reduction in the amount of adsorbed hydrocarbon and thus reduction in the amount of water and carbon dioxide generated in the regeneration unit.

- Reduction in unfavorable changes in the particle size distribution of the metal

- Improvement in the packing of a fixed bed reactor.

- Providing a hygroscopic metal halide to reactively remove water during upsets.

3. Starting the process with a solid which has undergone a number of regeneration cycles may offer benefits including:

- Reduction in byproducts.

- Improved chemical and thermal stability of reactors

4. Starting the process with the halogen present in part or completely as alkyl halide may be desirable for a number of reasons including:

- Less free halogen present during start-up.

- Ability to start the halide separations section early in the start-up sequence with no hydrogen halide or water present.

Shut-Down

1. Stopping the process with the halide in metal halide and/or alkyl halide form may improve safety, reduce corrosion, and improve maintenance accessibility.

2. Introducing reactive components into certain sections of the plant may provide a sink for halogen or hydrogen halide, improving safety, reducing corrosion and improving accessibility. An example of such a component is olefin.

Materials of Construction

1. The reactors for alkane halogenation and metathesis consisting of materials to minimize corrosion including but not limited to:

- Stainless steel;
- Silicon carbide;
- Glass lined steel;
- Titanium;
- Carbon fiber.

2. Process components operating at temperatures below 300 C. constructed from:

- Teflon;
- Glass.

What is claimed is:

1. A method comprising:

- providing a first quantity of alkane;
- providing a quantity of bromine;
- reacting the first quantity of alkane with the quantity of bromine to form alkyl monobromides, hydrogen bromide, and alkyl polybromides;
- providing a second quantity of alkane;
- reacting the second quantity of alkane with the alkyl polybromides to form additional alkyl monobromides;
- reacting the alkyl monobromides and the additional alkyl monobromides with a metal oxide to form a metal bromide and at least one reaction product selected from the group consisting of an alcohol and an ether;
- contacting the metal bromide with oxygen or air to regenerate the metal oxide and to form bromine; and
- recycling the bromine.

2. The method of claim 1 wherein the first and second quantities of alkane comprise at least one alkane selected from the group consisting of methane, ethane, propane, butane, and isobutane.

3. The method of claim 1 wherein the first and second quantities of alkane comprise methane.

13

4. The method of claim 1 wherein the alkyl polybromides comprise CH_2Br_2 , and the alkyl monobromides comprise CH_3Br .

5. The method of claim 1 wherein the reaction product comprises methanol.

6. The method of claim 1 wherein the reaction product comprises dimethyl ether.

7. The method of claim 1 wherein the alkyl monobromides and the additional alkylmonobromides are reacted with hydrogen bromide and the metal oxide to form the metal bromide and the at least one reaction product.

8. The method of claim 1 wherein the second quantity of alkane is reacted with the alkyl polybromides in the presence of an electrophilic catalyst.

9. A method comprising:

providing a first quantity of alkane;

providing a quantity of bromine;

reacting the first quantity of alkane with the quantity of bromine to form alkyl monobromides, hydrogen bromide, and alkyl polybromides;

providing a second quantity of alkane;

reacting the second quantity of alkane with the alkyl polybromides to form additional alkyl monobromides;

reacting the alkyl monobromides and the additional alkyl monobromides with a metal oxide to form a metal bromide and at least one reaction product selected from the group consisting of an alcohol, an ether, and an olefin; contacting the metal bromide with oxygen or air to regenerate the metal oxide and to form bromine; and recycling the bromine.

10. The method of claim 9 wherein the first and second quantities of alkane comprise at least one alkane selected from the group consisting of methane, ethane, propane, butane, isobutane, pentane, hexane, and cyclohexane.

11. The method of claim 9 wherein the first and second quantities of alkane comprise methane.

12. The method of claim 9 wherein the alkyl polybromides comprise at least one alkyl polybromide selected from the group consisting of CH_2Br_2 , CHBr_3 , and CBr_4 and wherein the alkyl monobromides comprise CH_3Br .

14

13. The method of claim 9 wherein the reaction product comprises methanol.

14. The method of claim 9 wherein the reaction product comprises dimethyl ether.

15. The method of claim 9 wherein the first and second quantities of alkane comprise two or more carbon atoms in their molecular structure and the reaction product comprises an olefin.

16. The method of claim 9 wherein the first and second quantities of alkane comprise ethane and the reaction product comprises at least one reaction product selected from the group consisting of diethyl ether, ethanol, and ethyl acetate.

17. The method of claim 9 wherein the alkyl monobromides and the additional alkylmonobromides are reacted with hydrogen bromide and the metal oxide to form the metal bromide and the at least one reaction product.

18. The method of claim 9 wherein the second quantity of alkane is reacted with the alkyl polybromides in the presence of an electrophilic catalyst.

19. A method comprising:

providing a first quantity of alkane;

providing a quantity of halogen;

reacting the first quantity of alkane with the quantity of halogen to form alkyl monohalides, hydrogen halide, and alkyl polyhalides;

providing a second quantity of alkane;

reacting the second quantity of alkane with the alkyl polyhalides to form additional alkyl monohalides;

reacting the alkyl monohalides and the additional alkyl monohalides with a metal oxide to form a metal halide and at least one reaction product selected from the group consisting of an alcohol, an ether, and an olefin;

contacting the metal halide with oxygen or air to regenerate the metal oxide and to form halogen; and recycling the halogen.

20. The method of claim 19 wherein the halogen comprises at least one halogen selected from the group consisting of chlorine, bromine, and iodine.

* * * * *